

# Four new species of *Ditrigona* Moore (Lepidoptera, Drepanidae) in China and an annotated catalogue

Xiao-Jiang Guo<sup>1,2</sup>, Rui Cheng<sup>1</sup>, Shan Jiang<sup>1</sup>, Da-Yong Xue<sup>1</sup>, Hong-Xiang Han<sup>1</sup>

**1** Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China **2** College of Life Science, Hebei University, Baoding, Hebei 071000, China

Corresponding author: Hongxiang Han ([hanhx@ioz.ac.cn](mailto:hanhx@ioz.ac.cn))

Academic editor: A. Hausmann | Received 7 December 2021 | Accepted 22 February 2022 | Published 31 March 2022

<http://zoobank.org/9A83F1CA-292E-41FC-A321-7B4719C51E7B>

**Citation:** Guo X-J, Cheng R, Jiang S, Xue D-Y, Han H-X (2022) Four new species of *Ditrigona* Moore (Lepidoptera, Drepanidae) in China and an annotated catalogue. ZooKeys 1091: 57–98. <https://doi.org/10.3897/zookeys.1091.78986>

## Abstract

The Chinese species of the genus *Ditrigona* Moore, 1888 are reviewed and an annotated catalogue is provided. Four new species are described from China: *Ditrigona sinespina* Jiang & Han, **sp. nov.**, *Ditrigona parva* Jiang & Han, **sp. nov.**, *Ditrigona concava* Guo & Han, **sp. nov.**, and *Ditrigona fusca* Guo & Han, **sp. nov.** *Derocra crystallina* Chu & Wang, 1987 and *Auzatella pentesticha* Chu & Wang, 1987 are newly combined into, respectively, the *derocina* and *quinaria* species groups of *Ditrigona*. *Ditrigona diana* Wilkinson is newly recorded in China. This results in 43 species of *Ditrigona* for the fauna of China. Illustrations of habitus and genitalia of the new species and most known species are presented.

## Keywords

DNA barcoding, Drepaninae, new combination, taxonomy

## Introduction

The genus *Ditrigona* was originally established by Moore (1888) on the basis of *Urapteryx triangularia* Moore, 1868 from India. Later, Warren (1922) described *Ditrigona regularis* Warren from Assam, and Bryk (1943) described *Ditrigona regularis differentiatia* Bryk from Burma. After a long silence in research into *Ditrigona*, Wilkinson (1968) provided the most comprehensive revision of the genus. He established three new generic synonyms of the genus, and transferred most of the species involved into *Ditrigona*; altogether he recorded 40 species and 12 subspecies for the genus, including

the description of 18 new species and five new subspecies. He also placed the species into four species groups, and provided descriptions and diagnosis of the genus, species groups, species, and subspecies. More recently, on the basis of Wilkinson's work, Chu and Wang (1988) recorded 36 species from China, including description of a new species *Ditrigona uniuncusa* Chu & Wang, and these species were included in vol. 3 of *Fauna Sinica* (Chu and Wang 1991). Holloway (1998) described two species *Ditrigona paludicola* and *Ditrigona wilkinsoni* from Borneo. The most recent research was provided by Li et al. (2015), who described *Ditrigona clavata* from Guangdong, China, and by Jiang and Han (2019), who described *Ditrigona tenuiata* from Sichuan, China and provided a checklist of the *triangularia* species group.

Further study of the specimens of *Ditrigona* from IZCAS and MHBUS shows that four new species need to be described. The purposes of this paper are to provide a survey and an annotated catalogue of Chinese *Ditrigona* species, to describe four new species, to newly record *Ditrigona diana* Wilkinson, 1968 from China, to transfer two species into the genus, and to provide illustrations of external features and genitalia of new species and most known species. This results in 43 species and 8 subspecies of *Ditrigona* for the fauna of China.

## Materials and methods

The depositories of all the types and examined specimens are indicated as follows:

<b>NHMUK</b>	The Natural History Museum, London, UK;
<b>ZFMK</b>	Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany;
<b>NHRS</b>	Naturhistoriska Riksmuseet, Stockholm, Sweden;
<b>IZCAS</b>	Institute of Zoology, Chinese Academy of Sciences, Beijing, China;
<b>MHBUS</b>	The Museum of Hebei University, Baoding, China;
<b>XTBG</b>	Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Yunnan, China;
<b>SCAU</b>	South China Agricultural University, Guangzhou, China;
<b>MNHN</b>	Muséum National d'Histoire Naturelle, Paris, France;
<b>DEI</b>	Deutsches Entomologisches Institut, Germany.

Terminology for the genitalia is based on Wilkinson (1968). Moths were photographed with a digital camera (Canon Pc1057). Composite images were generated using Auto-Montage software version 5.03.0061 (Synoptics Ltd). The sharpness-contrast of the photos was enhanced and the plates compiled using Adobe Photoshop (CS 5.1).

A total of 16 specimens of the species of the *triangularia* species group bearing a tail process were used for sequencing the DNA barcoding region of the mitochondrial COI gene. DNA barcodes of 15 specimens were successfully obtained in this work, and one sequence of *D. concava* was downloaded from BOLD: its related voucher specimen was donated by Prof. Akihior Nakamura from Xishuangbanna Tropical

**Table 1.** Details of specimens used in molecular analysis of the DNA barcode region.

Sample ID	Species	Date Collected	Locality	Collectors	GenBank/BOLD accession number
LEP M 33040	<i>D. parva</i>	6–8.Aug.2016	Tengchong, Yunnan	Ban XS	OL664050
LEP M 33049	<i>D. parva</i>	6–8.Aug.2016	Tengchong, Yunnan	Ban XS	OL664048
LEP M 33059	<i>D. parva</i>	6–8.Aug.2016	Tengchong, Yunnan	Ban XS	OL664049
LEP M 33016	<i>D. regularis</i>	9–12.Aug.2016	Yunlong, Yunnan	Ban XS	MK087682
LEP M 33027	<i>D. regularis</i>	6–8.Aug.2016	Tengchong, Yunnan	Ban XS	MK087683
LEP M 32911	<i>D. regularis</i>	10–13.Aug.2017	Xinping, Yunnan	Cui L	MK087678
LEP M 35671	<i>D. regularis</i>	14–16.Jul.2018	Anha, Sichuan	Cui L, Jiang S	MK087688
LEP M 32976	<i>D. triangularia</i>	13–14.Jul.2014	Weixi, Yunnan	Pan XD	MK087679
LEP M 25081	<i>D. tenuiata</i>	11.Sep.2016	Luding, Sichuan	Li XX	MK087687
LEP M 23038	<i>D. tenuiata</i>	7–10.Aug.2016	Kangding, Sichuan	Cui L	MK087685
LEP M 33029	<i>D. sinespina</i>	9–12.Aug.2016	Yunlong, Yunnan	Ban XS	MK087684
LEP M 33001	<i>D. sinespina</i>	9–12.Aug.2016	Yunlong, Yunnan	Ban XS	MK087680
LEP M 33002	<i>D. sinespina</i>	9–12.Aug.2016	Yunlong, Yunnan	Ban XS	MK087681
LEP M 33196	<i>D. sinespina</i>	26–27.Jun.2014	Tengchong, Yunnan	Pan XD	MK087677
LEP M 32975	<i>D. concava</i>	13–14.Jul.2014	Weixi, Yunnan	Li XX	OL664047
ARB00027811	<i>D. concava</i>	11.Aug.2011	Ailao Shan, Yunnan	Kitching RL, Ashton LA	SCDBC000200

Botanical Garden, Chinese Academy of Sciences (XTBG). Four of these specimens were *D. regularis*, one of *D. triangularia*, two *D. tenuiata*, four *D. sinespina* sp. nov., three *D. parva* sp. nov., and two *D. concava* sp. nov.

Protocols of DNA extraction and sequencing followed Ban et al. (2018). Details of studied specimens, including GenBank and BOLD accession numbers are summarized in Table 1. Pairwise distances within and between *Ditrigona* species for the COI barcoding region (612 bp) were calculated, and a neighbour-joining (NJ) tree (Saitou and Nei 1987) was constructed based on the Kimura two-parameter (K2P) method (Kimura 1980) using MEGA 6.0.

## Systematics

### Genus *Ditrigona* Moore, 1888

*Ditrigona* Moore, 1888: 258. Type species: *Urapteryx triangularia* Moore, 1867.

*Leucodrepana* Hampson, 1893: 333. Type species: *Leucodrepana idaeoides* Hampson, 1892.

*Leucodrepanilla* Strand, 1911: 198. Type species: *Corycia sacra* Butler, 1878.

*Auzatella* Strand, 1917: 148. Type species: *Auzata micronioides* Strand, 1917.

*Thaleridia* Moore, 1888: 266. Type species: *Thaleridia pruinosa* Moore, 1888.

**Generic characters.** The generic characters of *Ditrigona* and its differentiation from related genera are detailed in Wilkinson (1968) and Jiang and Han (2019).

**Distribution.** The species of *Ditrigona* are mainly distributed in the Oriental region.

### *derocina* species group

Wilkinson (1968) placed three species in the *derocina* species group: *Ditrigona derocina* (Bryk), *Ditrigona diana* Wilkinson, and *Ditrigona pruinosa* (Moore). The two former species have been recorded in China, and a further species, *Deroca crystallata* Chu & Wang, 1987 is newly placed in this species group in this work.

Species of the *derocina* species group are characterized by unipectinate antennae and semi-transparent wings. In the male genitalia, the *derocina* species group is unusual in *Ditrigona* in having sclerotization of the vinculum, and a quite long and narrow aedeagus. The eighth sternite is distinguished by large and curved octavals, and the tergite protrudes strongly. The female genitalia are characterized by having a very long ductus bursae, an ostial plate, an accessory sac and a long and narrow signum.

#### 1. *Ditrigona derocina* (Bryk, 1943)


Figs 1, 48, 82, 116, 152

*Peridrepana derocina* Bryk, 1943: 6. Holotype ♀, Burma: Kambaiti (NHRS).

*Ditrigona derocina*: Wilkinson, 1968: 418.

**Material examined.** **CHINA: Hubei** (IZCAS): 1♂7♀, Xuanen, Liangxihe, 796 m, 20–22.IX.2015, leg. Yao Jian, Zhao Kaidong. **Hunan** (IZCAS): 2♀, Sangzhi, Badagong Shan, Xiaozhuangping, 1420 m, 14.VI.2015, leg. Yao Jian, Zhao Kaidong. **Sichuan** (IZCAS): 1♀, Emei Shan, 0km, 1288 m, 31.VII.2013, leg. Cheng Rui. **Chongqing** (IZCAS): 1♂, Wu Shan, Wulipo, Dangyangcongping, 1773 m, 25.VII.2013, leg. Cheng Rui. **Yunnan** (IZCAS): 1♀, Lushui, Yaojiaping, 2500 m, 4.VI.1981, leg. Liao Subai; 2♂1♀, Tengchong, Houqiao, 1620 m, 6–8.VIII.2016, leg. Ban Xiaoshuang; 3♀, Tengchong, Heinitang, 1930 m, 28–30.V.1992, leg. Xue Dayong; 1♂1♀, same locality, 1824 m, 26–27.VI.2014, leg. Li Xinxin, Pan Xiaodan; 2♂3♀, Tengchong, Dahaoping, 2020 m, 24–26.V.1992, leg. Xue Dayong; 5♂2♀, same locality, 2020 m, 5–7.VIII.2007, leg. Wu Chunguang, Xue Dayong; 3♀, Lushui, Pianma, 1980 m, 3–4.VII.2014, leg. Pan Xiaodan; 1♀, Pianma, Dianxin hotel, 1970 m, 8–12.V.2011, leg. Yang Xiushuai, Wang Ke; 1♀, Gongshan, Puladi, 1298 m, 6–7.VII.2014, leg. Pan Xiaodan. **Tibet** (IZCAS): 1♂, Zham, 2400 m, 4.VII.1975, leg. Wang Ziqing; 2♀, same locality and collector, 2200 m, 23–30.VI.1957; 1♂1♀, Bomi, Tangmai, 2000 m, 26–28.VI.2015, leg. Li Xinxin; 1♀, Bomi, Tangmaidaqiao, 2037 m, 13–14.VI.2016, leg. Li Xinxin. **INDIA** (ZFMK): 1♂, Sikkim, Namchi, 1000 m, 2.VIII.1986, leg. W. Thomas, photograph examined.

**Distribution.** China (Hubei, Hunan, Sichuan, Chongqing, Yunnan, Tibet), India, Myanmar.



**Figures 1–27.** Adults of *Ditrigona* **1** *D. derocina*, male **2** *D. diana*, male **3** *D. crystallina*, holotype, male **4** *D. spilota*, male, ZFMK **5** *D. furvicosta*, male **6** *D. jordanaria*, male **7** *D. media*, paratype, male, ZFMK **8** *D. sericea*, male **9** *D. sericea*, male (*Auzatella pentesticha* Chu & Wang, allotype) **10** *D. pentesticha*, holotype, female **11** *D. q. erminea*, holotype, male, ZFMK **12** *D. q. spodia*, holotype, male, ZFMK **13** *D. q. leucophaea*, holotype, male, ZFMK **14** *D. obliquilinea thibetaria*, male **15** *D. triangularia*, lectotype, male **16** *D. uniuncusa*, male **17** *D. tenuiata*, holotype, male **18** *D. regularis*, male **19, 20** *D. sinespina* sp. nov., **19** holotype, male **20** paratype, female **21, 22** *D. parva* sp. nov., **21** holotype, male **22** paratype, female **23** *D. concava* sp. nov., holotype, male **24** *D. titana*, holotype, male, ZFMK **25** *D. pomenaria* male **26** *D. polyobotaria*, holotype, female, ZFMK **27** *D. typhodes*, male. Scale bar: 1 cm.

## 2. *Ditrigona diana* Wilkinson, 1968; new record for China

Figs 2, 49, 83, 117, 153

*Ditrigona diana* Wilkinson, 1968: 420. Holotype ♂, India: Gopaldara (NHMUK).

**Material examined.** **INDIA:** 1♂ (ZFMK), paratype, Khasis, IV.1895, Nat. Coll., Collectio. H.J. Elwes, moth photographed examined. **CHINA: Guangxi (IZCAS):** 1♂1♀, Napo, Defu, 1350 m, 19.VI.2000, leg. Li Wenzhu. **Yunnan (IZCAS):** 1♀, Xishuangbanna, Mengla, Menglun, 550 m, 12–15.V.2017, leg. Jiang Shan; 5♂2♀, Xishuangbanna, Mengla, Bubang, 680 m, 18–20.V.2017, leg. Jiang Shan; 1♂, Tengchong, Zhengding, 1833 m, 6–7.VIII.2013, leg. Li Xinxin; 1♂, Pingbian, Daweishan, 2090 m, 4–8.VIII.2017, leg. Cui Le; 1♂, Baoshan, Baihualing, 1520 m, 11–13.VIII.2007, leg. Wu Chunguang; 1♀, Tengchong, Dahaoping, 2020 m, 5–7.VIII.2007, leg. Wu Chunguang; 1♀, Tengchong, Heinitang, 1824 m, 26–27.VI.2014, leg. Pan Xiaodan; 1♀, Ruili, Wanding, Forest Garden, 900 m, 29.IV.2011, leg. Yang Xiushuai, Wang Ke; 2♀, Ruili, Wanding, Tianehu, 923 m, 30.IV.–1.V.2011, leg. Yang Xiushuai, Wang Ke; 8♀, Ruili, Mengmao, Mangling, 900 m, 26–27.IV.2011, leg. Yang Xiushuai, Wang Ke. **Tibet (IZCAS):** 1♂2♀, Mêdog, Yarang, 1091 m, 20–23.VIII.2006, leg. Lang Songyun.

**Distribution.** China (Guangxi, Yunnan, Tibet), India.

## 3. *Ditrigona crystallata* (Chu & Wang, 1987), comb. nov.

Figs 3, 50, 84, 118, 154

*Deroca crystallata* Chu & Wang, 1987: 116. Holotype ♂, China: Yunnan: Lushui: Yaojiaping (IZCAS).

**Note.** The species *Deroca crystallata* Chu & Wang, 1987 was described from Yunnan, Sichuan and Tibet. Its male genitalia obviously belong to the *derocina* species group of *Ditrigona*, and we therefore transfer the species to *Ditrigona*. The male genitalia are almost identical to those of *D. derocina*. However, the corpus bursae of the female genitalia is scobinate, which is different from the smooth ones of *D. derocina* and *D. diana*, though they share a very long and narrow ductus bursae and a slender curved signum. The ostial plate is invisible in *D. crystallata*, and it is most probably present, though it seems that the sternite was incorrectly removed and the abdomen is not preserved on the slide.

**Material examined.** **CHINA: Yunnan (IZCAS):** 1♂, holotype of *Deroca crystallata*, Lushui, Yaojiaping, 2500 m, 4.VI.1981, leg. Liao Subai; 1♂, Tengchong, Dahaoping, 2020 m, 24–26.V.1992, leg. Xue Dayong; 1♂, same locality, 5–7.VIII.2007, leg. Wu Chunguang, Xue Dayong. **Sichuan (IZCAS):** 1♀, allotype of *Deroca crystallata*, Guan Xian, Qingcheng Shan, 700–1600 m, 4.VI.1979, leg. Shang Jinwen. **Tibet (IZCAS):** 1♂, paratype of *Deroca crystallata*, Zham, 2400 m, 4.VII.1975, leg. Wang Ziqing.

**Distribution.** China (Sichuan, Yunnan, Tibet).

***quinaria* species group**

Wilkinson (1968) recorded 11 species in the *quinaria* species group: *Ditrigona spilota* Wilkinson, *Ditrigona inconspicua* (Leech), *Ditrigona furvicosta* (Hampson), *Ditrigona jordanaria* (Oberthür), *Ditrigona media* Wilkinson, *Ditrigona innotata* (Hampson), *Ditrigona sericea* (Leech), *Ditrigona quinaria* (Moore), *Ditrigona obliquilinea* (Hampson), *Ditrigona idaeoides* (Hampson), and *Ditrigona spatulata* Wilkinson. The former 10 species are recorded in China, and *Auzatella pentesticha* Chu & Wang is newly combined to *Ditrigona* in this work.

Species of the *quinaria* species group share bipectinate or serrate antennae with some species of the *triangularia* and *mytllata* species groups. In the male genitalia, the uncus is usually single, but sometimes bifurcate or notched. The single uncus resembles that of species of the *derocina* species group, but the group can be differentiated by the lack of sclerotization on the vinculum, and the large and broad saccus. The small and setose valva lacking a posterior projection also differs from those in the *triangularia* and *mytllata* species groups. The aedeagus is often characterized by the presence of a minute to large apical projection (not present in *D. spilota* and *D. obliquilinea*, and the situation unknown in *D. innotata* and *D. idaeoides*). The eighth sternite is modified with short octavals, and the tergite is often shallowly to moderately concave, occasionally straight or protruding with tiny lateral projections. The female genitalia lack an ostial plate; the ductus bursae is short and broad, and the corpus bursae usually has an accessory sac and a long and thin signum.

**4. *Ditrigona spilota* Wilkinson, 1968**

Figs 4, 51, 85, 119, 155

*Ditrigona spilota* Wilkinson, 1968: 423. Holotype ♂, China: Yunnan, Likiang (ZFMK).

**Material examined.** CHINA: Yunnan: 1♂ (ZFMK), paratype, Li-kiang (China), Provinz Nord-Yuennan, 16.VIII.1935, H. Höne, dissected in this work; 1♀ (ZFMK), same locality, 20.VIII.1935, H. Höne, dissected in this work; 1♀ (IZCAS), Tengchong, Danzhalinchang, 2500 m, 2–4.VI.1992, leg. Xue Dayong; 1♂1♀ (IZCAS), Lijiang, 3700 m, 9.VIII.2012, leg. Ashton; 2♂, same locality and collector, 3400 m, 17.VIII.2012. Sichuan (IZCAS): 1♀, Luding, Moxi, Hailuogou, 2596 m, 12.IX.2016, leg. Li Xinxin; 1♀, Luding, Moxi, Hailuogou Guancezhan, 3000 m, 10.IX.2016, leg. Li Xinxin.

**Distribution.** China (Sichuan, Yunnan).

**Remarks.** Compared to the male genitalia of the holotype (fig. 17 in Wilkinson 1968) of *D. spilota*, the socii of the paratype examined are much broader and blunter, while the aedeagus and 8<sup>th</sup> segment have no distinct differences. Further study is needed to investigate whether this is intraspecific variation, or more than one species is present in the large type series.

### 5. *Ditrigona inconspicua* (Leech, 1898)

*Teldenia inconspicua* Leech, 1898: 363. Lectotype ♂, China: Sichuan, Ta-Chien-lu (NHMUK).

*Peridrepana inconspicua*: Warren, 1922: 449.

*Ditrigona inconspicua*: Wilkinson, 1968: 425.

**Material examined.** No.

**Distribution.** China (Sichuan).

### 6. *Ditrigona furvicosta* (Hampson, 1912)

Figs 5, 52, 86, 120, 156

*Leucodrepana furvicosta* Hampson, 1912: 1271. Lectotype ♂, India: Sikkim (NHMUK).

*Ditrigona furvicosta*: Wilkinson, 1968: 428.

**Material examined.** CHINA: Yunnan: 1♀ (IZCAS), Tengchong, Danzhalinchang, 2500 m, 2–4.VI.1992, leg. Xue Dayong; 1♂1♀ (IZCAS), same locality, 2479 m, 30.VI.–1.VII.2014, leg. Pan Xiaodan; 2♂ (IZCAS), Lijiang, Alpine Botanical Garden, 3260–3452 m, 20.VI.2009, leg. Qi Feng; 1♂ (ZFMK), Li-kiang (China), Provinz Nord-Yuennan, 23.VI.1935, H. Höne, moth photograph examined. Tibet (IZCAS): 2♀, Yadong, Yadonglinchang, 2690 m, 24.VI.2016, leg. Li Xinxin; 1♂3♀, Nyingchi, Pêlung, 2115 m, 1.IX.2005, leg. Wang Xuejian; 1♂ (MHBU), Zham, 27.VII.2005, leg. Shi Aimin.

**Distribution.** China (Yunnan, Tibet), India.

### 7. *Ditrigona jordanaria* (Oberthür, 1923)

Figs 6, 53, 87, 121

*Corycia jordanaria* Oberthür, 1923: 238. Lectotype ♂, China: Sichuan, Ta-t sien-lu (ZFMK).

*Ditrigona jordanaria*: Wilkinson, 1968: 429.

**Material examined.** CHINA: Henan (IZCAS): 1♂, Baiyun Shan, 1550 m, 13–15.VIII.2008, leg. Xue Dayong, Song Wenhui. Shaanxi (IZCAS): 1♂, Ningshan, Huoditang, 1520 m, 13–17.VIII.2016, leg. Cheng Rui, Jiang Shan. Sichuan: 1♂ (IZCAS), Luding, Hailuogou, 3010 m, 2–4.VIII.2014, leg. Pan Xiaodan; 1♂ (ZFMK), Ta-t sien-Lou, 1899, Chasseurs indigenes, Ex. Oberthür Coll., Brit. Mus. 1927-3, moth photograph examined.

**Distribution.** China (Henan, Shaanxi, Sichuan, Tibet).

**8. *Ditrigona media* Wilkinson, 1968**

Figs 7, 54, 88, 122

*Ditrigona media* Wilkinson, 1968: 431. Holotype ♂, China: Sichuan, Ta-tsien-Lou (NHMUK).

**Material examined.** CHINA: **Sichuan** (ZFMK): 1♂, paratype, Batang, Tibet [Sichuan], Alpine Zone, ca. 5000 m, 7.VI.1936, H. Höne, moth photograph examined; 1♂, same locality, 23.VI.1938, H. Höne, dissected in this work. **Gansu** (IZCAS): 1♂, Dangchang, Guanegou, 2045 m, 1–3.VIII.2016, leg. Cheng Rui, Jiang Shan.

**Distribution.** China (Gansu, Sichuan, Tibet).

**9. *Ditrigona innotata* (Hampson, 1893)**

*Drepana innotata* Hampson, 1893: 335. Lectotype ♀, China: Kulu (Young) (NHMUK).

*Peridrepana innotata*: Warren, 1922: 449.

*Ditrigona innotata*: Wilkinson, 1968: 433.

**Material examined.** No.

**Distribution.** China (Tibet).

**10. *Ditrigona sericea* (Leech, 1898)**

Figs 8–9, 55–56, 89–90, 123, 157

*Teldenia sericea* Leech, 1898: 263. Lectotype ♂, China: Sichuan, Moupin (NHMUK).

*Drepana fulvicosta* Dudgeon, 1899: 652. Syntype, India.

*Peridrepana fulvicosta*: Gaede, 1931: 7.

*Leucodrepana nivea brimanica* Bryk, 1943: 7. Holotype ♀ (as ♂): Burma: Kambaiti (NHRS).

*Ditrigona sericea*: Wilkinson, 1968: 434.

**Material examined.** CHINA: **Shaanxi** (IZCAS): 2♀, Ningshan, Huoditang, 1520 m, 13–17.VIII.2016, leg. Cheng Rui, Jiang Shan. **Sichuan**: 3♂1♀ (IZCAS), Luding, Moxi, Boyangcun, 1691 m, 1.VIII.2014, leg. Li Xinxin; 1♀ (IZCAS), Baoxing, Dashuigou, Guanhuozhan, 1591 m, 1–5.VIII.2016, leg. Cui Le; 1♀ (IZCAS), Emei Shan, Qingyinge, 800–1000 m, 14.V.1957, leg. Zhu Fuxing; 1♂ (ZFMK), Ta-Tsien-Lou, Tche To, Chasseurs Indigènes, 1894, Ex. Oberthür Coll., Brit. Mus. 1927-3, moth photograph examined; 1♂ (ZFMK), Siao-Lou, 1903, Coll. R. P. Déjean. **Yunnan** (IZCAS): 1♂1♀, Tengchong, Houqiao, 1553 m, 28–29.VI.2014, Pan Xiaodan, Li Xinxin; 11♂17♀,

same locality, 1620 m, 6–8.VIII.2016, leg. Ban Xiaoshuang; 1♀, same locality, 1080 m, 31.V.–1.VI.1992, leg. Xue Dayong; 3♂5♀, Tengchong, Dahaoping, 2020 m, 24–26.V.1992, leg. Xue Dayong; 3♂1♀, same locality, 2020 m, 5–7.VIII.2007, leg. Xue Dayong; 5♂1♀, same locality and date, leg. Wu Chunguang; 1♂2♀, ibidem, leg. Lang Songyun; 1♂2♀, Tengchong, Heinitang, 1824 m, 26–27.VI.2014, leg. Pan Xiaodan; 1♀, same locality and date, leg. Li Xinxin; 1♂, Tengchong Shidi, 1730 m, 3–5.VIII.2016, leg. Ban Xiaoshuang; 1♂3♀, Tengchong, Qushi, Dabacun, 1873 m, 4.VIII.2013, leg. Liu Shuxian; 2♂, same locality, 1823 m, 5.VIII.2013, leg. Liu Shuxian; 15♂5♀, Gongshan, Dulongjiang, 1505 m, 8–9.VII.2014, leg. Pan Xiaodan; 1♀, Gaoligong, Nankang, 2000 m, 21.III.2007, leg. Zhang Peiyi; 1♀, Gaoligong, Baihualing, 1500 m, 16.IX.2007, leg. Zhang Peiyi; 1♂, Baoshan, Bawan, 1040 m, 8–10.VIII.2007, leg. Wu Chunguang; 1♂, Pingbian, Dawei Shan, 2043 m, 19–20.VII.2016, leg. Ban Xiaoshuang; 5♂7♀, same locality, 2090 m, 4–8.VIII.2017, leg. Cui Le; 1♀, Yongsheng, Liude, 2300 m, 9.VII.1984, leg. Chen Yixin; 3♂1♀, Kunming, Xishan, 2100 m, 23.III.1958, leg. Meng Xuwu; 1♂, Lushui, Pianma, 1980 m, 3–4.VII.2014, leg. Li Xinxin; 1♂, Weixi, Pantiange, 2570 m, 15–16.VII.2014, leg. Pan Xiaodan; 2♂, Dali, Cangshan, 2226 m, 23–24.VI.2014, leg. Pan Xiaodan; 3♂1♀, same locality and date, leg. Li Xinxin; 1♂1♀, Yunlong, Tianchi, 2570 m, 9–12.VIII.2016, leg. Ban Xiaoshuang; 1♂2♀, Yunlong, Tianchi, 2570 m, 9–12.VIII.2016, leg. Ban Xiaoshuang; 2♂, Ailao Shan, 2000 m, 19–20.VIII.2011, leg. Ashton, ex. XTBG. **TIBET** (IZCAS): 1♂, allotype of *Auzatella pentesticha* Chu & Wang, 1987, Zham, 2400 m, 26.VI.1975, leg. Huang Fusheng; 1♂, Bomi, Tangmai, 2000 m, 26–28.VI.2015, leg. Li Xinxin; 2♂7♀, Mêdog, Hanmi, 2095 m, 10–11.VIII.2006, leg. Lang Songyun; 1♂3♀, Mêdog, Aniqiao, 1060 m, 12–13.VIII.2006, leg. Lang Songyun; 1♀, Mêdog, Dayandong, 2880 m, 9.VIII.2006, leg. Lang Songyun; 1♀, Mêdog 108K, 848 m, 4.VIII.2014, leg. Cheng Rui, Cui Le; 1♂2♀, Nyingchi, Pêlung, 1900 m, 24–25.VI.2015, leg. Li Xinxin; 1♂1♀, same locality, 2115 m, 1.IX.2005, leg. Wang Xuejian; 1♀, Nyingchi, Pêlung, Mamba, 2115 m, 1–2.IX.2005, leg. Wang Xuejian; 1♂1♀, Cona, Lexiang, Senmuzha, 2741 m, 2–3.VI.2016, leg. Li Xinxin; 1♀, Cona, Lexiang, Lewangdaqiao, 2423 m, 7.VI.2016, leg. Li Xinxin; 1♂, Yadong, Yadonglinchang, 2690 m, 24.VI.2016, leg. Li Xinxin; 1♂, Gyirong, Resuo, 18.VIII.1984, leg. Pu Qiongqiong; 1♀, Gyirong, Tuowu, 3300 m, 4.VIII.1975, leg. Huang Fusheng; 1♀, Zham, Kouan, 26.IX.1984, leg. Li Aihua; 1♀, Zham, 20.IX.1984, leg. Guo Sengbao, 1♀, Zham, Daqu, 3300 m, 2.VII.1957, leg. Wang Ziqing; 1♀, Zham, Nyalam, 2200 m, 9.V.1966, leg. Wang Shuyong. **MYANMAR** (ZFMK): 1♀, Upper Burma Htawgaw, 6000ft, Coll. A.E. Swann.

**Distribution.** China (Shaanxi, Sichuan, Yunnan, Tibet), India, Myanmar.

**Remarks.** Chu and Wang (1987) described *Auzatella pentesticha* based on four specimens from Tibet and Hubei Province, and designated the female from Quxam, Tibet as the holotype, the male from Zham, Tibet as the allotype, and another two females from Hubei as paratypes. Unfortunately, the genitalia slide of the female holotype could not be found. In the original description, the figure of the female genitalia is from one of the paratypes from Hubei Province. The only male from Tibet has genitalia identical to those of *D. sericea*, and the two females from Hubei belong to *D. quinaria*. Although the holotype and allotype specimens were collected from two very close localities (Quxam

and Zham are less than 30 km apart), and in a very similar season (7 July, 26 June), we hesitate to synonymize *Auzatella pentesticha* Chu & Wang with *Ditrigona sericea* (Leech) without having seen the genitalia of the holotype; however, we redetermine the male allotype as *D. sericea*, and the two paratypes as *Ditrigona quinaria erminea* Wilkinson.

**11. *Ditrigona pentesticha* (Chu & Wang, 1987), comb. nov.**

Fig. 10

*Auzatella pentesticha* Chu & Wang, 1987: 108. Holotype ♀, China: Tibet: Quxam (IZCAS).

**Material examined.** CHINA: Tibet (IZCAS): 1 ♀, holotype, Quxam, 3300 m, 7.VII.1975, leg. Wang Ziqing.

**Distribution.** China (Tibet).

**Remarks.** As stated under the above species, this species now only includes the female holotype. The validity of the species needs further study, for example, by obtaining a DNA barcode from the holotype.

**12. *Ditrigona quinaria* (Moore, 1867)**

*Drepanodes quinaria* Moore, 1867: 618. Neotype ♂, India: Darjiling (NHMUK).

*Ditrigona quinaria*: Wilkinson, 1968: 438.

**Note.** At present, *D. quinaria* comprises five subspecies; four are recorded from China, the exception being *D. quinaria nivea* (Hampson), which is distributed in India.

***Ditrigona quinaria quinaria* (Moore, 1867)**

**Material examined.** No.

**Distribution.** China (Tibet), India.

***Ditrigona quinaria erminea* Wilkinson, 1968**

Figs 11, 57, 91, 124, 158

*Ditrigona quinaria erminea* Wilkinson, 1968: 442. Holotype ♂, China: Shaanxi, Tapaishan-im-Tsinling (ZFMK).

**Material examined.** CHINA: Shaanxi (ZFMK): 1 ♂, holotype, Tapaishan im Tsinling, Sued-Shensi, ca. 3000 m, 26.VI.1936, H. Höne, slide no. 1493, moth photograph examined; 1 ♂, Tapaishan im Tsinling, Sued-Shensi, China, ca. 3000 m, 11.VIII.1936,

H. Höne, dissected in this work; 1♀, same locality and collector, 17.VI.1936, dissected in this work. **Hubei** (IZCAS): 2♀, paratypes of *Auzatella pentesticha* Chu & Wang, 1987, Shennongjia, Juhulinchang, 1840 m, 16.VIII.1981, leg. Han Yinheng.

**Distribution.** China (Shaanxi, Hubei).

### *Ditrigona quinaria spodia* Wilkinson, 1968

Figs 12, 58, 92, 125

*Ditrigona quinaria spodia* Wilkinson, 1968: 442. Holotype ♂, China: Yunnan, A-tun-tse (ZFMK).

**Material examined.** CHINA: **Yunnan:** 1♂ (ZFMK), holotype, A-tun-tse (N Yunnan), Aus Höhe ca. 4000 m, 25.VII.1937, H. Höne, Drepanidae genitalia slide No. 1487, moth photograph examined; 1♀ (ZFMK), Paratype, A-tun-tse (N. Yunnan), Aus Höhe, ca. 4000 m, 15.VI.1937, H. Höne; 1♂ (ZFMK), paratype, Li-kiang, ca. 3000 m, Prov. Nord-Yuennan, 15.IV.1934, H. Höne; 1♂1♀ (IZCAS), Tengchong, Heinitang, 1930 m, 28–30.V.1992, leg. Xue Dayong; 1♂ (IZCAS), Xianggelila, Gezan, 3141 m, 20–21.VII.2014, leg. Li Xinxin; 1♂ (IZCAS), Weixi, Tacheng, 2800 m, 13–14.VII.2014, leg. Li Xinxin.

**Distribution.** China (Yunnan).

### *Ditrigona quinaria leucophaea* Wilkinson, 1968

Fig. 13

*Ditrigona quinaria leucophaea* Wilkinson, 1968: 443. Holotype ♂, China: Tibet [Sichuan], Batang (ZFMK).

**Material examined.** CHINA: **Sichuan:** 1♂ (ZFMK), holotype, Batang (Tibet), Im Tal dea Yantze, ca. 2800 m, 16.V.1936, H. Höne, Drepanidae genitalia slide No. 1486, moth photograph examined.


**Distribution.** China (Sichuan).

## 13. *Ditrigona obliquilinea* (Hampson, 1893)

*Leucodrepana obliquilinea* Hampson, 1893: 333. Lectotype ♂, India: Assam, Naga Hills (NHMUK).

*Ditrigona obliquilinea:* Wilkinson, 1968: 444.

**Note.** *Ditrigona obliquilinea* includes two subspecies, and the nominate subspecies is distributed in India and Myanmar.



**Figures 28–47.** Adults of *Ditrigona* **28** *D. lineata lineata*, male **29** *D. lineata tephroides*, holotype, male, ZFMK **30** *D. legnichrysa*, paratype, male, ZFMK **31** *D. policharia*, lectotype, female, ZFMK **32** *D. artema*, male **33** *D. candida*, paratype, male, ZFMK **34** *D. chionea*, paratype, male, ZFMK **35**, **36** *D. fusca* sp. nov., holotype, male **35** upperside **36** underside **37–38** *D. conflexaria micronioides* **37** male **38** female **39** *D. conflexaria cerodeta*, holotype, male, ZFMK **40** *D. margarita*, male **41** *D. berres*, male **42** *D. chama*, male **43** *D. platytes*, male **44** *D. clavata*, male **45** *D. marmorea*, paratype, female ZFMK **46** *D. aphyta*, male **47** *D. cirruncata*, male. Scale bar: 1 cm.

***Ditrigona obliquilinea thibetaria* (Poujade, 1895)**

Figs 14, 59, 93, 126, 159

*Micronia thibetaria* Poujade, 1895: 311. Lectotype ♂, China: Thibet [Sichuan], Moupin (MNHN).

*Leucodrepana thibetaria*: Leech, 1898: 311.

*Corycia pnocaria* Oberthür, 1923: 238

*Ditrigona obliquilinea thibetaria*: Wilkinson, 1968: 445.

**Material examined.** CHINA: Hunan (IZCAS): 1♂4♀, Sangzhi, Badagong Shan, Xiaozhuangping, 1420 m, 14.VI.2015, leg. Yao Jian, Zhao Kaidong. Sichuan: 2♂ (IZCAS), Jiguan Shan, Shaoyaogou, 1556 m, 11–16.VII.2016, leg. Cui Le; 1♂ (ZFMK), Tien-Tsuen, Yui-Kin, 1899, Chasseurs indigènes, moth photograph examined. Tibet (IZCAS): 1♀, Gyirong, 26.V.1984, leg. Daci; 1♂, Cona, Mama, 2930 m, 18–20.VI.2015, leg. Li Xinxin; 1♂, Cona, Lexiang, Senmuzha, 2741 m, 2–3.VI.2016, leg. Li Xinxin.

**Distribution.** China (Shaanxi, Zhejiang, Hubei, Hunan, Sichuan, Tibet).

**14. *Ditrigona idaeoides* (Hampson, 1893)**

*Leucodrepana idaeoides* Hampson, 1893: 333. Lectotype ♂, Sikkim: Tonglo (NHMUK).

*Ditrigona idaeoides*: Wilkinson, 1968: 447.

**Material examined.** No

**Distribution.** China (Sichuan), India.

***triangularia* species group**

Based on Wilkinson (1968) and Jiang and Han (2019), the *triangularia* species group contains ten species, in which *Ditrigona triangularia* (Moore), *Ditrigona regularis* Warren, *Ditrigona uniuncusa* Chu & Wang, and *Ditrigona tenuiata* Jiang & Han bear elongate posterior projections of the hind wings (tail process); the other six species, *Ditrigona titana* Wilkinson, *Ditrigona pomenaria* (Oberthür), *Ditrigona typhodes* Wilkinson, *Ditrigona polybotaria* (Oberthür), *Ditrigona sciara* Wilkinson, and *Ditrigona fasciata* (Hampson) lack the tail process. The first nine species are recorded in China, and three new species (*D. sinespina*, *D. parva*, *D. concava*) with the tail process are described in this work.

The species with a tail process on the hind wing have quite distinct wing patterns: the hind wing has the postmedial and submarginal lines approaching each other near the anal angle, and bears a small black patch at the upper angle of the tail. The species lacking a tail process resemble some species of the *mytylata* species group, in that they have transverse lines which often resemble a narrow band. In the male genitalia, the valva is characterized by having a small flap-like extension. The species with a tail

process can also be distinguished by the large rounded socii and the stout aedeagus bearing a brush-like cornutus. In the species lacking a tail process, the aedeagus is narrow, straight or bent, and the cornutus is a simple process or absent. The eighth sternite is small, shallowly concave or protruding in species with a tail process, and the eighth tergite almost unmodified. Both eighth tergite and sternite often possess octavals in the species lacking a tail process. In the female genitalia, the ostium bursae is usually large, and the ductus bursae is often indiscernible, but wide and obvious in *D. typhodes*. The corpus bursae bears a small accessory sac in species with a tail process. (modified from Wilkinson, 1968)

16 DAN barcoding sequences were obtained for *D. regularis*, *D. triangularia*, *D. tenuiata*, *D. concava* sp. nov., *D. parva* sp. nov., and *D. sinespina* sp. nov., and the six species are clearly separated from each other in the COI barcode fragment (fig. 178). The genetic distance between these species is 8.92% (min. 7.16%, max. 12.32%).

### 15. *Ditrigona triangularia* (Moore, 1867)

Figs 15, 60, 94, 127

*Urapteryx triangularia* Moore, 1867: 612. Lectotype ♂, India: Darjiling (NHMUK).  
*Ditrigona triangularia*: Moore, 1888: 258.

**Material examined.** INDIA: 1♂ (NHMUK), lectotype, Darjiling, Moore Coll. 94-106, moth photograph examined. CHINA: Yunnan: 1♂ (IZCAS), Weixi, Tacheng, 2800 m, 13–14.VII.2014, leg. Pan Xiaodan; 1♂ (ZFMK), Li-kiang (China), Provinz Nord-Yuennan, 28.VI.1935, H. Höne, moth photograph examined.

**Distribution.** China (Fujian, Taiwan, Yunnan, Sichuan), India, Myanmar.


### 16. *Ditrigona uniuncusa* Chu & Wang, 1988

Figs 16, 61, 95, 128, 160

*Ditrigona uniuncusa* Chu & Wang, 1988: 202. Holotype ♂, China: Fujian, Wuyi Shan (IZCAS).

**Material examined.** CHINA: Fujian (IZCAS): 1♂, holotype, Wuyi Shan, 22.VI.1982, leg. Zhang Baolin; 1♀, same locality, 704 m, 12.VIII.1979, leg. Song Shimei; 1♀, Wuyi Shan, Sangang, 704 m, 23.X.1980, leg. Cai Rongquan; 1♀, same locality, 1.VI.1983, leg. Mai Guoqing; 3♀, same locality, 31.VII.2005, leg. Wang Jiashe; 1♂20♀, same locality, 20–21.X.2005, leg. Han Hongxiang, Lang Songyun, Yang Chao; 2♀, same locality, X.1979, leg. Huang Juyi; 1♀, same locality, 8.X.1979, leg. Xu Zhanfei; 1♀, same locality, 15.VI.1981, leg. Jiang Fan; 1♀, same locality, 21.VI.1981, leg. Wang Jiashe, Jiang Fan. Sichuan (IZCAS): 1♀, Luding, Moxi, 19–20.V.2009, leg. Li Jing.

**Distribution.** China (Fujian, Sichuan).



**Figures 48–63.** Male genitalia of *Ditrigona* **48** *D. derocina* **49** *D. diana* **50** *D. crystallina*, holotype **51** *D. spilota*, ZFMK **52** *D. furvicosta* **53** *D. jordanaria* **54** *D. media* **55** *D. sericea* **56** *D. sericea* (*Auzatella pentesticha* Chu & Wang, allotype) **57** *D. quinaria erminea*, ZFMK **58** *D. quinaria spodia* **59** *D. obliquilinea tibetaria* **60** *D. triangularia* **61** *D. uniuncusa* **62** *D. tenuiata*, holotype **63** *D. regularis*. Scale bars: 1 mm.

**17. *Ditrigona tenuiata* Jiang & Han, 2019**

Figs 17, 62, 96, 129, 161

*Ditrigona tenuiata* Jiang & Han, 2019: 84. Holotype, ♂, China: Sichuan, Kangding (IZCAS).

**Material examined.** CHINA: Sichuan (IZCAS): 1♂, holotype, Kangding, Xikangyinxiang hotel, 2582 m, 7–10.VIII.2016, leg. Cui Le, M23028; 1♀, paratype, Luding, Hailuoguo, 2569 m, 11.IX.2016, leg. Li Xinxin, M25081.

**Distribution.** China (Sichuan).

**18. *Ditrigona regularis* Warren, 1922**

Figs 18, 63, 97, 130, 162

*Ditrigona regularis* Warren, 1922: 463. Lectotype ♂, India: Assam, Khasia (NHMUK).  
*Ditrigona regularis difflerentiata* Bryk, 1943: 9.

**Material examined.** CHINA: **Guangxi** (IZCAS): 2♀, Napo, Defu, 1350 m, 19.VI.2000, leg. Li Wenzhu. Sichuan: 1♀, Emei Shan, Qingyinge, 800–1000 m, 29.IV.1957, leg. Huang Keren; 2♀, Emei Shan, 0km, 1288 m, 31.VII.2013, leg. Cheng Rui; 1♀, Huili, 19.VII.1974, leg. Han Yinheng; 2♂1♀, Anha, Luoji Shan, 2044 m, 14–16.VII.2018, leg. Cui Le, Jiang Shan. **Yunnan** (IZCAS): 1♀, Pianma, Dianxin hotel, 1970 m, 8–12.V.2011, leg. Yang Xiushuai, Wang Ke; 1♀, Tengchong, Dahaoping, 2020 m, 24–26.V.1992, leg. Xue Dayong; 1♂, same locality, 2020 m, 5–7.VIII.2007, leg. Xue Dayong; 1♀, Tengchong, Shidi, 1730 m, 3–5.VIII.2016, leg. Ban Xiaoshuang; 2♂2♀, Tengchong, Houqiao, 1620 m, 6–8.VIII.2016, leg. Ban Xiaoshuang; 1♀, Lushui, Yaojiaping, 2500 m, 4.VI.1981, leg. Zhang Xuezhong; 1♀, Ruili, Dengga, 980 m, 6–8.VI.1992, leg. Xue Dayong; 2♂3♀, Pingbian, Daweishan, 2090 m, 4–8.VIII.2017, leg. Cui Le; 1♀, Kunming, Shuanglongxiang, 2100 m, 11.VIII.2006, leg. Ma Rong; 1♀, Xinping, Gasa, Yaonan, 1900 m, 10–13.VIII.2017, leg. Cui Le; 1♀, Yunlong, Tianchi Baohuqu, 2570 m, 9–12.VIII.2016, leg. Ban Xiaoshuang. **Tibet** (IZCAS): 1♀, Medôg, 1091 m, 22.VIII.2006, leg. Lang Songyun. **THAILAND** (ZFMK): 1♂, Chiangmai Doi Suthep, 1325 m, 21.XI.–4.XII.1989, leg. Schnitzler, moth photograph examined.

**Distribution.** China (Guangxi, Sichuan, Yunnan, Tibet), India, Myanmar, Thailand.

**19. *Ditrigona sinespina* Jiang & Han, sp. nov.**

<http://zoobank.org/B95C0171-9D11-419C-B668-371D07B000E3>

Figs 19–20, 64, 98, 131, 163

**Description. Head.** Antennae bipectinate, with proximal rami shorter than outer rami, the longest ramus about four times diameter of antennal shaft in male; rami quite short in female, almost equal to diameter of antennal shaft. Frons flattened, width less than

diameter of compound eyes; white, upper half with a narrow pale brown transverse band. Labial palpus slender, not extending beyond frons, with outside brown, inner side whitish. Vertex white, pale brown anteriorly.

**Thorax.** Dorsal and ventral sides of thorax white. Tegula white. Hind tibia with two pairs of spurs in both sexes. Forewing length: ♂♀ 16 mm. Both fore- and hind wings white, transverse lines grey. Forewing with costa pale brown, distal half deeper. Sub-basal and antemedial lines slightly bent inwards at middle and costa, the former narrower; postmedial line broad, almost straight; submarginal line double, the inner one slightly wavy and the outer one deeply wavy. Hind wing with antemedial line straight, merging into the elongate grey area along anal margin; postmedial line broader, almost straight, closing to submarginal line near anal angle, forming large pointed teeth on CuA<sub>2</sub> and anal fold; submarginal line double, with the inner one nearly straight and only wavy near anal angle, the outer one wavy, the two lines gradually approximating towards anal margin. Anal margin less extended, possessing a quite short tail process, longer in female, with a small black patch. Fringes pale brown. Forewing underside with costa deep brown in basal half.

**Abdomen.** Dorsal and ventral sides of abdomen white. Eighth tergite large, nearly quadrate, with posterior margin shallowly concave; eighth sternite concave at middle, forming two small lateral blunt processes.

**Male genitalia.** Uncus bifurcate over its whole length, both halves short and very narrow. Socii large, rounded. Valva small, ventral margin smoothly curved, distal and posterior margins straight, forming a blunt angle; posterior protrusion rounded. Saccus blunt and rounded. Juxta indistinctly shaped. Aedeagus very stout, terminal part narrower; cornutus a large oval spinose patch.

**Female genitalia.** Papillae analis short; apophyses anteriores moderate, broad basally. Ostium bursae large; ductus bursae indiscernible; corpus bursae round, signum absent.

**Diagnosis.** On the wing pattern, *D. sinespina* is close to *D. tenuiata*, but it can be differentiated by the larger distance between the two submarginal lines on the forewing, and the smaller tail process. Compared to *D. triangularia* and *D. uniuncusa*, the anal margin of *D. sinespina* is less extended, and the tail process is distinctly shorter than in those two species. Compared to *D. parva* sp. nov. and *D. concava* sp. nov., *D. sinespina* is larger (with forewing length 16 mm), and the tail process on the hind wing is less developed.

In the male genitalia, the slender uncus is similar to that of *D. tenuiata* and *D. concava*, but it is longer than in *D. tenuiata* and shorter than in *D. concava*. The straight distal margin of the valva is also different from these two species. The shape of the aedeagus, which is broad and blunt posteriorly, also can be distinguished from these two species. The eighth tergite of the male is similar to that of *D. concava*, but the eighth sternite is different: in *D. sinespina* it is narrowly and deeply concave, forming two blunt protrusions, while in *D. concava* it is widely and shallowly concave, forming two small lateral processes. The female genitalia of *D. sinespina* are also similar to those of *D. tenuiata*, but can be differentiated by the lack of a signum.

**Type material.** Holotype, ♂, CHINA: Yunnan (IZCAS): Yunlong, Tianchi, 2570 m, 9–12.VIII.2016, leg. Ban Xiaoshuang, slide no. Drep-1054, M33001. Paratypes: Yunnan (IZCAS), 1♂, same data as holotype, M33002, posterior part of abdomen missing; 1♂, same data as holotype, M33029; 1♀, Tengchong, Heinitang, 1824 m, 26–27.VI.2014, leg. Pan Xiaodan, slide no. Drep-1060, M33196.

**Distribution.** China (Yunnan).

**Etymology.** The species is named from the Latin words *sine* and *spina*, which refers to the lack of a signum in the female genitalia.

**Molecular data.** The mean intraspecific distance of *D. sinespina* is 1.55% (min. 0%, max. 2.24%,  $n = 4$ ). The nearest related species is *D. tenuiata*, with genetic distance 7.16%.

## 20. *Ditrigona parva* Jiang & Han, sp. nov.

<http://zoobank.org/3E199CD5-F712-4F67-B30E-E04C94EA34E9>

Figs 21–22, 65, 99, 132, 164

**Description.** Head and thorax almost identical to those of *D. sinespina*. Forewing length: ♂11.5 mm, ♀14 mm. Antemedial line almost straight apart from an inward bend at costa. Outer line of the double submarginal lines serrate. Anal margin of hind wing elongate, with relatively large tail process. Fringes brown. Forewing underside with costa brown at basal half.

**Male genitalia.** Uncus bifurcate over whole length, both sides broad. Socii large, terminally semicircular, scobinate. Valva small, distal margin shallowly concave, posterior protrusion rounded. Saccus blunt and rounded. Juxta with posterior margin almost straight. Aedeagus very stout; cornutus a large oval spinose patch. Eighth tergite quadrate, with a pair of small anterior apodemes; eighth sternite quite small, posterior margin slightly convex.

**Female genitalia.** Papillae analis short; apophyses anteriores moderate, broad basally. Ostium bursae large; ductus bursae indiscernible; corpus bursae rounded, posteriorly with a large wrinkled sclerotized area and accessory sac, signum short and narrow.

**Diagnosis.** The wing pattern is very close to that of *D. uniuncusa*. The antemedial line on the hind wing is straight in *D. parva*, but slightly convex in *D. uniuncusa*. The width between the two submarginal lines is larger than in *D. uniuncusa*, especially in the female. In the male genitalia, *D. parva* shares a stout uncus with *D. uniuncusa* and *D. regularis*, but the terminal half of the socii are quite different: scobinate and semicircular in *D. parva*, scobinate and tapering in *D. uniuncusa*, and spinose in *D. regularis*. The female genitalia are also different: the sclerotized area on the corpus bursae is rounded, less sclerotized and smaller than that in *D. uniuncusa*, which has a large oval well sclerotized area; the signum is shorter than in *D. uniuncusa*.

**Type material.** Holotype, ♂, CHINA: Yunnan (IZCAS): Tengchong, Houqiao, 1620 m, 6–8.VIII.2016, leg. Ban Xiaoshuang, slide no. Drep-1057, M33049.

Paratypes: 1♂, same data as holotype, M33059; 1♀, same data as holotype, slide no. Drep-1059, M33040; 1♂, same locality, 1553 m, 28–29.VI.2014, leg. Pan Xiaodan.

**Distribution.** China (Yunnan).

**Etymology.** The species is named referring to the Latin word *parvus*, which refers to the small wings.

**Molecular data.** The three specimens of *D. parva* have no genetic distance between them. The nearest related species is *D. concava*, with a genetic distance of 9.56%.

## 21. *Ditrigona concava* Guo & Han, sp. nov.

<http://zoobank.org/AB4E8887-3422-4197-AA1C-DC8363354992>

Figs 23, 66, 100, 133

**Description.** Characters of head and thorax in male same as in *D. sinespina*. Forewing length 16 mm in male. Fore- and hind wings white, transverse lines grey. Forewing with costa pale brown. Subbasal line slightly bent inwards at middle and costa; antemedial line straight and only bent inwards near costa; postmedial line broad, almost straight; submarginal line double, the inner one slightly wavy and the outer one deeply wavy. Hind wing with antemedial line slightly convex at middle, bent outwards and merging into the elongate grey area along the anal margin; postmedial line broader, almost straight, slanting outwards and closing to meet submarginal line near anal angle, forming large pointed teeth on  $CuA_2$  and anal fold; submarginal line double, with the inner one nearly straight and only wavy near anal angle, the outer one wavy, the two lines gradually approximating towards anal margin. Anal margin less extended, with a quite short tail process, longer in the female, with a small black patch. Fringes pale brown. Forewing underside with costa deep brown in basal half.


**Male genitalia.** Uncus bifurcate over its whole length, both arms slender. Socii large, terminally semicircular, rough apically. Valva small, distal and posterior margin shallowly concave, posterior protrusion rounded. Saccus blunt and rounded. Juxta large, slightly sclerotized, indistinctly shaped. Aedeagus stout, terminal part narrow; cornutus an oblong spinose patch. Eighth tergite quadrate, posterior margin almost straight; eighth sternite shallowly concave, with two small lateral protrusions.

**Female genitalia.** Unknown.

**Diagnosis.** The most distinctive character of *D. concava* lies in the male eighth sternite, which is different from all other congeners by the wide and shallow concavity, bearing two small lateral processes. The male genitalia are similar to those of *D. tenuiata* and *D. sinespina*, and can be differentiated by the following differences: the uncus is longer than in those two species; the distal and posterior margins of the valva are shallowly concave in *D. concava*, but the two margins are straight in *D. sinespina*.

**Type material.** Holotype, ♂, **CHINA: Yunnan** (IZCAS): Ailao Shan, 2600 m, 11.VIII.2011, leg. Kitching Ashton, slide no. Drep-1095, ARB00027811, ex. XTBG. Paratype: 1♂, **Yunnan** (IZCAS): Weixi, Tacheng, 2800 m, 13–14.VII.2014, leg. Li Xinxin, slide no. Drep-1091, M32975.

**Distribution.** China (Yunnan).



**Figures 64–74.** Male genitalia of *Ditrigona* **64** *D. sinespina* sp. nov., holotype **65** *D. parva* sp. nov., holotype **66** *D. concava* sp. nov., paratype **67** *D. titana*, ZFMK **68** *D. pomenaria* **69** *D. typhodes* **70** *D. lineata lineata* **71** *D. artema* **72** *D. candida*, paratype, ZFMK **73** *D. chionea* **74** *D. fusca* sp. nov., holotype. Scale bars: 1 mm.

**Etymology.** The species is named after the Latin word *concauus*, which refers to the shallowly concave 8<sup>th</sup> male sternite.

**Molecular data.** An intraspecific distance of *D. sinespina* of 2.07% ( $n = 2$ ) was recorded. The nearest related species is *D. parva*, with a genetic distance of 9.56%.

## 22. *Ditrigona titana* Wilkinson, 1968

Figs 24, 67, 101, 134

*Ditrigona titana* Wilkinson, 1968: 453. Holotype ♂, China: Yunnan, Likiang (ZFMK).

**Material examined.** CHINA: Yunnan: 1♂ (ZFMK), holotype, Li-kiang, ca. 3000 m, Prov. Nord-Yuennan, 13.IX.1934, H. Höne, moth photograph examined.

**Distribution.** China (Yunnan).

## 23. *Ditrigona sciara* Wilkinson, 1968

*Ditrigona sciara* Wilkinson, 1968: 458. Holotype ♂, China: Sichuan, Ta-tsien-lou (NHMUK).

**Material examined.** No.

**Distribution.** China (Sichuan).

## 24. *Ditrigona pomenaria* (Oberthür, 1923)

Figs 25, 68, 102, 135

*Corycia (Bapta) pomenaria* Oberthür, 1923: 238. Lectotype ♂, China: Sichuan, Moupin (ZFMK).

*Ditrigona pornenaria*: Wilkinson, 1968: 454.

**Material examined.** CHINA: Sichuan (ZFMK): 1♂, holotype, Mou-Pin, 1897, ex. R.P. Déjean, photograph examined; 1♂, Jiguan Shan, Shaoyaogou, 1556 m, 11–16. VII.2016, leg. Cui Le.

**Distribution.** China (Sichuan).

## 25. *Ditrigona polyobotaria* (Oberthür, 1923)

Fig. 26

*Corycia polyobotaria* Oberthür, 1923: 237. Lectotype ♀, China: Sichuan, Siao-lou (ZFMK).

*Ditrigona polyobotaria*: Wilkinson, 1968: 458.

**Material examined.** CHINA: **Sichuan** (ZFMK): 1♀, holotype, Siao-Lou, 1900, Chasseurs indigènes, photograph examined.

**Distribution.** China (Sichuan).

## 26. *Ditrigona typhodes* Wilkinson, 1968

Figs 27, 69, 103, 136, 165

*Ditrigona typhodes* Wilkinson, 1968: 456. Holotype ♂, China: Yunnan, Likiang (ZFMK).


**Material examined.** CHINA: **Yunnan**: 1♂ (ZFMK), holotype, Li-kiang, ca. 3000 m, Prov. Nord-Yuennan, 19.VIII.1934, H. Höne, moth photograph examined; 5♂1♀ (IZCAS), Ailao Shan, 2400–2600 m, 6–13.VIII.2011, leg. Ashton, ex. XTBG; 4♂4♀ (IZCAS), Lijiang, 3200–3600 m, 9–14.VIII.2012, leg. Ashton; 2♀ (IZCAS), Lijiang, Yulongshan, 22.VII.–2.VIII.1962, leg. Song Shimei; 1♂ (IZCAS), Lijiang Alpine Botanic Garden, 3272 m, 15–16.VIII.2013, leg. Li Xinxin; 1♀ (IZCAS), Weixi, Tacheng, 2800 m, 13–14.VII.2014, leg. Pan Xiaodan; 1♀ (IZCAS), Lushui, Yaojiaping, 2500 m, 4.VI.1981, leg. Liao Subai; 2♀, Yongsheng, Liude, 2250 m, 10.VII.1984, leg. Liu Dajun. **Sichuan** (IZCAS): 1♂2♀, Yajiang, Bingzhan, 3340 m, 30–31.VII.2014, leg. Li Xinxin.

**Distribution.** China (Sichuan, Yunnan), Myanmar.

## *mytilata* species group

In Wilkinson (1968) and Li et al. (2015), 15 species in the *mytilata* species group were recorded in China: *Ditrigona lineata* (Leech), *Ditrigona legnichrysa* Wilkinson, *Ditrigona policharia* (Oberthür), *Ditrigona artema* Wilkinson, *Ditrigona marmorea* Wilkinson, *Ditrigona candida* Wilkinson, *Ditrigona conflexaria* (Walker), *Ditrigona margarita* Wilkinson, *Ditrigona aphyra* Wilkinson, *Ditrigona berres* Wilkinson, *Ditrigona chama* Wilkinson, *Ditrigona chionea* Wilkinson, *Ditrigona platytes* Wilkinson, *Ditrigona cirruncata* Wilkinson, and *Ditrigona clavata* Li & Wang. A new species *D. fusca* is described in this work.

This species group is characterized by usually having lamellate antennae, and the valva of the male genitalia usually possessing a long posterior extension. Other characters are summarized as follows: the forewing is sometimes weakly falcate; the streaks vary considerably, with transverse lines linear, band-like or absent; the uncus is bifurcate, and the socii are very large; the aedeagus is usually arcuate; both eighth sternite and tergite are modified, the former with short octavals, and the latter concave with small or large protrusions; in the female genitalia, the ostial pocket is characteristic, and the corpus bursae lacks an accessory sac. (modified from Wilkinson, 1968)



**Figures 75–89.** (75–81) Male genitalia of *Ditrigona* **75** *D. conflexaria micronioides* **76** *D. margarita* **77** *D. berres* **78** *D. chama* **79** *D. platytes* **80** *D. clavata* **81** *D. cirruncata*. (82–89) Aedeagus of *Ditrigona* **82** *D. derocina* **83** *D. diana* **84** *D. crystallata*, holotype **85** *D. spilota*, ZFMK **86** *D. furvicosta* **87** *D. jordanaria* **88** *D. media* **89** *D. sericea*. Scale bars: 1 mm.

**27. *Ditrigona lineata* (Leech, 1898)**

*Leucodrepana lineata* Leech, 1898: 364. Holotype ♂, China: Sichuan, Omei-Shan (NHMUK).

*Ditrigona lineata*: Wilkinson, 1968: 462.

Currently, *D. lineata* includes two subspecies, and both are distributed in China.

***Ditrigona lineata lineata* (Leech, 1898)**

Figs 28, 70, 104, 137, 166

**Material examined.** CHINA: Sichuan (IZCAS): 1♂, Emei Shan, 15.VIII.1977; 1♂1♀, Emei Shan, Leidongping, 2444 m, 7–8.VIII.2014, leg. Pan Xiaodan; 2♀, Pingwu, Wanglang, Changbaigou, 2480 m, 24.VII.2016, leg. Cui Le. Yunnan: 2♂ (IZCAS), Lijiang, Yulong Shan, 30.VII., 4.VIII.1962, leg. Song Shimei; 1♂ (ZFMK), Li-kiang, China, 6.VIII.1935, Coll. H. Höne.

**Distribution.** China (Sichuan, Yunnan).

***Ditrigona lineata tephroides* Wilkinson, 1968**

Fig. 29

*Ditrigona lineata tephroides* Wilkinson, 1968: 464. Holotype ♂, China: Shaanxi, Tapaishan-im-Tsinling (ZFMK).

**Material examined.** CHINA: Shaanxi (ZFMK): 1♂, holotype, Tapaishan im Tsinling, Sued-Shensi, ca. 3000 m, 12.VIII.1936, H. Höne, moth and genitalia photos examined.

**Distribution.** China (Shaanxi, Tibet).

**28. *Ditrigona legnichrysa* Wilkinson, 1968**

Fig. 30

*Ditrigona legnichrysa* Wilkinson, 1968: 466. Holotype ♂, China: Tibet [Sichuan] (NHMUK).

**Material examined.** CHINA: Sichuan (ZFMK): 1♂, paratype, Tien-Tsuen, 1897, ex. R.P. Déjean, moth photograph examined.

**Distribution.** China (Zhejiang, Sichuan, Yunnan, Tibet).

**29. *Ditrigona policharia* (Oberthür, 1923)**

Fig. 31

*Corycia (Bapta) policharia* Oberthür, 1923: 237. Lectotype ♀, China: Sichuan, Tsien-Tsuen (ZFMK).

*Ditrigona policharia*: Wilkinson, 1968: 468.

**Material examined.** CHINA: Sichuan (ZFMK): 1 ♀, lectotype, Tien-Tsuen, 1897, ex. R. P. Déjean, moth photograph examined.

**Distribution.** China (Sichuan).

**Remarks.** This species was described based on a single female specimen only. Wilkinson (1968) stated that the female genitalia are similar to those of *Ditrigona legnichrysa* Wilkinson, and he also had difficulty in distinguishing it from *D. legnichrysa*, though he listed several tiny differences. He suggested that further material was needed to clarify the relationship between the two species.

**30. *Ditrigona artema* Wilkinson, 1968**

Figs 32, 71, 105, 138, 167

*Ditrigona artema* Wilkinson, 1968: 469. Holotype ♂, China: Sichuan, Siao-lou (NHMUK).

**Material examined.** CHINA: Sichuan: 1 ♂ (ZFMK), paratype, Ta-tsien-Lou, Chasseurs du P. Déjean, 1904, Ex. Oberthür Coll., Brit. Mus. 1927-3, moth photograph examined; 1 ♀ (ZFMK), paratype, Wahuipass, 4000 m, Süd. Tatsienlu, VII.–VIII.1930, leg. Friedrich, Coll. Dr. Wehril, moth photograph examined; 1 ♀ (IZCAS), Emei Shan, 15.VIII.1977; 2 ♂ (IZCAS), Luding, Hailuogou Erhaoyingdi, 2569 m, 11.IX.2016, leg. Li Xinxin; 1 ♂6 ♀ (IZCAS), Luding, Moxi, Hailuogou, 2596 m, 12.IX.2016, leg. Li Xinxin. Tibet (IZCAS): 1 ♂, Yadong, 2760 m, 23–25.VIII.2014, leg. Cheng Rui, Cui Le.

**Distribution.** China (Sichuan, Tibet).

**31. *Ditrigona candida* Wilkinson, 1968**

Figs 33, 72, 106, 139, 168

*Ditrigona candida* Wilkinson, 1968: 472. Holotype ♂, China: Yunnan, Likiang (ZFMK).

**Material examined.** CHINA: Yunnan (ZFMK): 1 ♂, holotype, Li-kiang (China), Provinz Nord-Yuennan, 5.VIII.1935, H. Höne, moth photograph examined; 1 ♂, paratype, Li-kiang, ca. 2000 m, Prov. Nord-Yuennan, 15.VII.1934, H. Höne, dissected in this work; 1 ♀, paratype, Li-kiang (China), Provinz Nord-Yuennan, 2.VIII.1935, H. Höne, dissected in this work.

**Distribution.** China (Yunnan).

**32. *Ditrigona chionea* Wilkinson, 1968**

Figs 34, 73, 107, 140, 169

*Ditrigona chionea* Wilkinson, 1968: 490. Holotype ♂, China: 'Chasseurs Thibetains' (NHMUK).

**Material examined.** CHINA: Yunnan: 1♂1♀ (ZFMK), paratypes, Li-kiang. ca. 3000 m, Prov. Nord-Yuennan, 24.VII.1934, 14.VI.1934, H. Höne, moth photograph examined; 1♂6♀ (IZCAS), Lijiang Alpine Botanic Garden, 3260–3452 m, 15–20.VI.2009, leg. Xue Dayong, Yang Chao, Han Hongxiang, Qi Feng; 1♀ (IZCAS), Lijiang, Wenhai, 3097 m, 19.VI.2009, leg. Xue Dayong; 1♀ (IZCAS), Lijiang, Ganheba, 3296 m, 23.VI.2009, leg. Qi Feng. Shaanxi (IZCAS): 1♀, Zhouzhi, Diaoyutai, 1480 m, 29.VI.2008, leg. Bai Ming.

**Distribution.** China (Shaanxi, Hubei, Sichuan, Yunnan).

**33. *Ditrigona fusca* Guo & Han, sp. nov.**

<http://zoobank.org/ED43239E-A79D-4818-B3D3-D4F58984B228>

Figs 35–36, 74, 108, 141, 170

**Description.** *Head.* Antennae simple in both male and female. Frons yellow, width less than diameter of compound eyes. Labial palpus with outside deep brown, inner side yellowish. Vertex pale yellow.

*Thorax.* Tegula pale brown. Dorsal and ventral sides of thorax pale brown. Hind tibia with two pairs of spurs in both sexes. Forewing length: ♂16 mm, ♀15–17 mm. Both wings pale brown, evenly decorated with brown scales, less in basal half of hind wing. Transverse lines absent. Fringes yellowish brown. Underside with forewing deep brown, distal part paler, costa yellowish brown; hind wing with costa yellowish brown, other parts identical to upperside. Fringes yellowish brown.

*Abdomen.* Dorsal and ventral sides of abdomen pale brown. The eighth tergite with posterior margin concave, with two blunt lateral protrusions; the eighth sternite with posterior margin slightly convex, with two small hooked lateral processes.

*Male genitalia.* Uncus bifurcate over whole length, the arms narrow and slender. Socii large, tongue-like, of even width, with tips blunt and scobinate. Valva nearly triangular, with tip blunt, ventral margin decorated with tiny spines; basal posterior process bent, tip expanded, with a small accompanying bursa. Juxta rounded, with posterior margin protruding, mound-like. Saccus narrow. Aedeagus slender, almost even in width, tip blunt.

*Female genitalia.* Ostial pocket band-like. Lamella antevaginalis paired leaf-like. Ductus bursae indiscernible. Corpus bursae rounded; signum a narrow longitudinal sclerotized strip; accessory sac absent.

**Diagnosis.** The wing pattern is distinctive in lacking transverse lines on both fore- and hind wings. The male genitalia are very close to those of *D. candida*, *D. chionea* and *D. margarita* in the *mytylata* species group, in that they share the slender bifid uncus and tongue-like socii. *D. fusca* and *D. chionea* can be differentiated from those two species by

the broader valva, on the base of which a sclerotized ridge is present. The difference between *D. fusca* and *D. chionea* in the male genitalia lies in the shape of the juxta, which is widely protruding posteriorly in *D. fusca*, but only with a tiny process at middle in *D. chionea*. The aedeagus is also different, straight and almost even in width in *D. fusca*, but tapering and twisted in *D. chionea*. The female genitalia of *D. fusca* and *D. chionea* are almost identical.

**Type material.** Holotype, ♂ (IZCAS), **CHINA: Yunnan:** Lijiang, Yulong Shan, 23.VIII.1962, leg. Song Shimei, slide no. Drep-1092. Paratypes (IZCAS): **Yunnan:** same locality and collector as holotype, 1♂, 3.VIII.1962, slide no. Drep-954; 1♀, 30.VII.1962; 2♀, 2.VIII.1962; 1♀, 3.VIII.1962, 2900 m; 3♀, 4.VIII.1962, slide no. Drep-953; 1♀, 23.VIII.1962; 1♀, 25.VIII.1962; 4♀, 30.VIII.1962; 1♀, Xianggelila, Xiaozhongdian, 3235 m, 15–16.VIII.2016, leg. Ban Xiaoshuang, slide no. Drep-1093; 1♀, Lijiang, Alpine Botanical Garden, 3260–3452 m, 16–18.VI.2009, leg. Qi Feng.

**Distribution.** China (Yunnan).

**Etymology.** The specific name is from the Latin word *fuscus*, which refers to the pale brown wing colour.

### 34. *Ditrigona conflexaria* (Walker, 1861)

*Acidalia conflexaria* Walker, 1861: 148. Holotype ♂, N. China (NHMUK).

*Ditrigona conflexaria*: Wilkinson, 1968: 475.

Based on Wilkinson (1968), *D. conflexaria* comprises three subspecies, all distributed in China.

### *Ditrigona conflexaria conflexaria* (Walker, 1861)

**Material examined.** No.

**Distribution.** North China.

### *Ditrigona conflexaria micronioides* (Strand, 1917)

Figs 37–38, 75, 109, 142, 171


*Auzata (Auzatella) micronioides* Strand, 1917: 148. Holotype ♀, China: Formosa (DEI).

*Leucodrepana micronioides*: Watson, 1959: 232.

*Auzatella micronioides*: Inoue, 1962: 12.

*Ditrigona conflexaria micronioides*: Wilkinson, 1968: 475.

**Material examined. CHINA (IZCAS): Shanxi:** 1♂1♀, Yicheng, Dahelinchang, 1212 m, 13–15.VIII.2018, leg. Zhang Xinyi; 1♀, Yuanqu, Huangguman, 1258 m, 21–22.VIII.2018, leg. Zhang Xinyi. **Henan:** 1♂1♀, Baiyun Shan, 1550 m, 13–15.VIII.2008, leg. Song Wenhui; 1♂1♀, Nanyang, Baotianman, 14.VII.2006, 27.VII.2006, leg.



**Figures 90–106.** Aedeagus of *Ditrigona* **90** *D. sericea* (*Auzatella pentesticha* Chu & Wang, allotype) **91** *D. quinaria erminea*, ZFMK **92** *D. quinaria spodia* **93** *D. obliquilinea thibetaria* **94** *D. triangularia* **95** *D. uniuncusa* **96** *D. tenuiata*, holotype **97** *D. regularis* **98** *D. sinespina* sp. nov., holotype **99** *D. parva* sp. nov., holotype **100** *D. concava* sp. nov., paratype **101** *D. titana*, ZFMK **102** *D. pomenaria* **103** *D. typhodes* **104** *D. lineata lineata* **105** *D. artema* **106** *D. candida*, ZFMK. Scale bars: 1 mm.

Shen Xiaocheng et al.; 13♂1♀, Baotianman, 1407 m, 10–11.VIII.2008, leg. Jiang Nan, Song Wenhui, Xue Dayong. **Shaanxi**: 1♂, Shangnan, Jinsixia, 777 m, 23–25.VII.2013, leg. Jiang Nan; 5♂, same locality, 766 m, 16–19.VII.2017, leg. Cui Le; 3♂, Ningshan, Yueba, 1052 m, 1–3.VIII.2018, leg. Zhang Xinyi; 3♂, Mei Xian, Honghegu, Shenxianling, 1239 m, 21–22.VII.2018, leg. Zhang Xinyi; 2♀, same locality, 874 m, 20.VII.2018, leg. Zhang Xinyi; 6♂, Foping, Longcaoping, 1218 m, 4.VIII.2018, leg. Zhang Xinyi; 4♂, Taibai, Huangbaiyuan, 1279 m, 15–17.VII.2018, leg. Zhang Xinyi. **Zhejiang**: 5♂, Zhoushan, Putuo, Taohuadiao, 40 m, 4.VIII.2016, leg. Li Xinxin; 1♂, Taishun, Wuyanling, Shangfengxiang, 1050 m, 30.VII.2005, leg. Lang Songyun. **Hubei**: 8♂4♀, Lichuan, Xingdoushan, Sanxianchang, 1144 m, 17–19.V.2017, leg. Li Henan; 5♂, Xuanen, Changtanhe, Lianghekou, 949 m, 13–14.V.2017, leg. Li Henan; 5♂2♀, Xuanen, Changtanhe, Dawolong, 713–794 m, 15–16.V.2017, leg. Li Henan; 1♀, Ying Shan, Wujiashan, 500 m, 30.VI.2014, leg. Xue Dayong. **Jiangxi**: 1♂1♀, Jinggangshan, Huangyangjie, 1090 m, 4.VIII.2013, leg. Pan Xiaodan. **Hunan**: 3♀, Zhangjiajie, Wulingyuan, Wenfeng, 475 m, 10.VI.2015, leg. Yao Jian, Zhao Kaidong; 1♂1♀, Yanling, Taoyuandong, 631 m, 4–8.VII.2008, leg. Chen Fuqiang. **Fujian**: 1♂, Meihua Shan, Huyuan, 1276 m, 19.VII.2013, leg. Xue Dayong; 1♀, Masu, 25.IX.1981, leg. Jiang Fan. **Guangxi**: 1♀, Maoershan, Jiuniuchang, 1150 m, 7.VII.1985, leg. Fang Chenglai; 1♀, Maoershan, Jiuniutang, 1146 m, 16.VIII.2012, leg. Yang Chao; 1♂, Huanjiang, Yangmeiao, 1189 m, 18–22.VII.2015, leg. Jiang Nan. **Sichuan**: 1♂, Anha, Luojuoshan, 2044 m, 14–16.VII.2018, leg. Cui Le, Jiang Shan; 1♂, Hongya, Wawushan, Jinhuaqiao, 1147 m, 12–14.VIII.2016, leg. Cui Le; 12♂6♀, Emei Shan, Qingyinge, 800–1000 m, 16.IV., 17.IV., 19.IV., 24.IV., 26.IV., 27.IV., 29.IV., 30.IV., 12.V., 17.V., 18.VII.1957, leg. Huang Keren, Zhu Fuxing, Lu Youcai, Wang Zongyuan; 1♂, Emei Shan, 1100 m, 22.VI.1955, leg. Li Jinhua; 2♂, Wan Xian, Wangerbao, 1200 m, 27.IX.1994, leg. Song Shimei; 1♂ (ZFMK), Ost Tien-mu-shan, Chekiang, 14.VII.1931, H. Höne, moth photograph examined. **Chongqing**: 1♀, Jinyun Shan, 800 m, 13.VI.1994, leg. Li Wenzhu; 2♂4♀, Chongqing, 800 m, 20.VI., 22.VI.1974, leg. Han Yinheng. **Guizhou**: 2♀, Qianping Shan, Fudingshan, 604 m, 1–4.V.2018, leg. Zhao Kaidong.

**Distribution.** China (Shanxi, Henan, Shaanxi, Zhejiang, Hubei, Jiangxi, Hunan, Fujian, Taiwan, Guangxi, Sichuan, Chongqing, Guizhou), Japan.

### *Ditrigona conflexaria cerodeta* Wilkinson, 1968

Fig. 39


*Ditrigona conflexaria cerodeta* Wilkinson, 1968: 477. Holotype ♂, China: Likang (ZFMK).

**Material examined.** CHINA: **Yunnan**: 1♂ (ZFMK), holotype, Li-kiang, ca. 3000 m, Prov. Nord-Yuennan, 8.VIII.1934, H. Höne, moth photograph examined; 1♀ (ZFMK), paratype, Li-kiang, ca. 3000 m, 4.VIII.1934, H. Höne, moth photograph examined.

**Distribution.** China (Yunnan).



**Figures 107–127.** (107–115) Aedeagus of *Ditrigona* **107** *D. chionea* **108** *D. fusca* sp. nov., holotype **109** *D. confluxaria micronioides* **110** *D. margarita* **111** *D. berres* **112** *D. chama* **113** *D. platytes* **114** *D. clavata* **115** *D. cirruncata*. Scale bars: 1 mm. (116–127) Eighth segment of *Ditrigona* **116** *D. derocina* **117** *D. diana* **118** *D. crystallina*, holotype **119** *D. spilota*, ZFMK **120** *D. furvicosta* **121** *D. jordanaria* **122** *D. media* **123** *D. sericea* **124** *D. quinaria erminea*, ZFMK **125** *D. quinaria spodia* **126** *D. obliquilinea thibetaria* **127** *D. triangularia*



**Figures 128–151.** Eighth segment of *Ditrigona* **128** *D. uniuncusa* **129** *D. tenuiata*, holotype **130** *D. regularis* **131** *D. sinespina* sp. nov., holotype **132** *D. parva* sp. nov., holotype **133** *D. concava*, paratype **134** *D. titana*, ZFMK **135** *D. pomenaria* **136** *D. typhodes* **137** *D. lineata lineata* **138** *D. artema* **139** *D. candida*, ZFMK **140** *D. chionea* **141** *D. fusca*, sp. nov., holotype **142** *D. conflexaria micronioides* **143** *D. margarita* **144** *D. berres* **145** *D. chama* **146** *D. platytes* **147** *D. clavata* **148–151** *D. cirruncata* **148** from Shaanxi **149–150** from Emei Shan, Sichuan **151** from Zhejiang.

**35. *Ditrigona margarita* Wilkinson, 1968**

Figs 40, 76, 110, 143, 172

*Ditrigona margarita* Wilkinson, 1968: 483. Holotype ♂, China: Shaanxi, Tapaishan-im-Tsinling (ZFMK).

**Material examined. CHINA: Shaanxi:** 1♂ (ZFMK), holotype, Tapaishan-im-Tsinling, Sued-Shensi. ca. 1700 m, 22.VI.1936, H. Höne, moth photograph examined; 1♂ (IZCAS), Ningshan, Huoditang, 1538 m, 11–15.VII.2012, leg. Cheng Rui; 5♂ (IZCAS), Nanzheng, Liping, 1540 m, 27–30.VII.2017, leg. Li Henan. **Henan** (IZCAS): 1♂, Baotianman, 27.VII.2006, leg. Shen Xiaocheng et al. **Ningxia** (IZCAS): 1♂, Jingyuan, Qiuqianjialinchang, 1822 m, 23.VI.2008, leg. Song Wenhui; 1♀, Jingyuan, Erlonghe, 1984 m, 11–12.VII.2008, leg. Song Wenhui. **Gansu** (IZCAS): 1♀, Kang Xian, Qinghelinchang, 1400 m, 8.VII.1999, leg. Zhu Chaodong; 1♀, Zhouqu, Shatanlinchang, 2400 m, 14.VII.1999, leg. Yao Jian; 1♀, Wen Xian, Qiujiaba, 2350 m, 21.VII.1999, leg. Zhu Chaodong; 1♀, same locality and date, leg. Yao Jian. **Sichuan** (IZCAS): 1♂, Mao Xian, 9–11.VII.2015, leg. Li Xinxin; 1♀, Jiguan Shan, Baliping, 1470 m, 15–16.VII.2016, leg. Cui Le; 1♂, Jiguan Shan, Shaoyaogou, 1556 m, 11–16.VII.2016, leg. Cui Le; 2♀, Pingwu, Wanglang, Qikeshu, 2446 m, 21–22.VII.2016, leg. Cui Le.

**Distribution.** China (Shanxi, Henan, Shaanxi, Ningxia, Gansu, Sichuan).

**36. *Ditrigona berres* Wilkinson, 1968**

Figs 41, 77, 111, 144, 173

*Ditrigona berres* Wilkinson, 1968: 486. Holotype ♂, China: Shaanxi, Tapaishan-im-Tsinling (ZFMK).

**Material examined. CHINA: Shaanxi:** 1♂ (ZFMK), holotype, Tapaishan-im-Tsinling Sued-Shensi, ca. 3000 m, 23.VI.1936, H. Höne, moth photograph examined; 1♂ (IZCAS), Feng Xian, Jialingjiangyuantou, 1510 m, 21–24.VII.2017, leg. Cui Le. **Hubei** (IZCAS): 2♀, Shennongjia, Dajiuhu, 1800 m, 1.VIII.1981, leg. Han Yinheng; 1♂, Xingshan, Longmenhe, 1300 m, 14.VI.1993, leg. Li Hongxing. **Hunan** (IZCAS): 2♂1♀, Sangzhi, Badagongshan, Xiaozhuangping, 1420 m, 18.VI.2015, leg. Yao Jian, Zhao Kaidong.

**Distribution.** China (Shaanxi, Hubei, Hunan).

**37. *Ditrigona chama* Wilkinson, 1968**

Figs 42, 78, 112, 145, 174

*Ditrigona chama* Wilkinson, 1968: 488. Holotype ♂, China: Sichuan, Siao-lou (NHMUK).

**Material examined. CHINA: Yunnan:** 1♂ (ZFMK), paratype, Li-kiang. ca. 3000 m, Prov. Nord-Yuennan, 28.VII.1934, H. Höne, moth photograph examined; 2♂

(IZCAS), Yongsheng, Liude, 2250 m, 10.VII.1984, leg. Liu Dajun; 1♂ (IZCAS), Tengchong, Qushi, Dabacun, 1873 m, 4.VII.2013, leg. Liu Shuxian. **Shanxi** (IZCAS): 15♂14♀, Pangquangou, Erhezhuang, 1670 m, 4–6.VII.2018, leg. Cui Le, Jiang Shan. **Shaanxi** (IZCAS): 1♂, Mei Xian, Honghegu, Shenxianling, 1239 m, 21–22.VII.2018, leg. Zhang Xinyi; 1♂, Ningshan, Huoditang, 1497 m, 29–31.VII.2018, leg. Zhang Xinyi; 1♀, Zhouzhi, Diaoyutai, 1480 m, 29.VI.2008, leg. Bai Ming; 1♂, Foping, Longcaoping, 1256 m, 3.VII.2008, leg. Liu Wangang, Cui Junzhi; 2♂, Taibai, Huangbaiyuan, 1323 m, 17–18.VI.2012, leg. Li Jing, Liu Shuxian; 1♂1♀, Feng Xian, Jialingjiangyantou, 1510 m, 21–24.VII.2017, leg. Cui Le; 1♂1♀, Zhashui, Yingpanzhen, Niubeiliang, Laolin, 1046 m, 11–15.VII.2017, leg. Cui Le. **Gansu** (IZCAS): 1♂, Kang Xian, Qinghelinchang, 1400 m, 7.VII.1999, leg. He Tongli, Yao Jian; 1♂, Zhouqu, Shatanlinchang, 2350 m, 5.VII.1998, leg. Yuan Decheng. **Zhejiang**: 2♀, Tianmu Shan, 26.VI.1957, leg. Su Jiyao. **Sichuan** (IZCAS): 1♀, Emei Shan, Jiulaodong, 1800–1900 m, 6.VII.1957, leg. Zhu Fuxing; 1♀, Jiguan Shan, Shaoyaogou, 1556 m, 11–16.VII.2016, leg. Cui Le.

**Distribution.** China (Shanxi, Shaanxi, Gansu, Zhejiang, Hubei, Sichuan, Yunnan, Tibet).


### 38. *Ditrigona platytes* Wilkinson, 1968

Figs 43, 79, 113, 146, 175

*Ditrigona platytes* Wilkinson, 1968: 492. Holotype ♂, China: Chekiang, West Tien-mu-Shan (ZFMK).

**Material examined.** **CHINA: Zhejiang:** 1♂ (ZFMK), holotype, West Tien-mu-shan, Prov. Chekiang, 29.V.1932, H. Höne, moth photograph examined; 1♂ (IZCAS), Tianmu Shan, 26.VI.1957, leg. Su Jiyao; 2♂ (IZCAS), West Tianmu Shan, Qianmutian, 1330 m, 30.VII.2011, leg. Yan Keji, Cheng Rui. **Fujian:** 1♀ (ZFMK), paratype, Kuantun, 2300 m, 26.V.1938, leg. J. Klapperich, moth photograph examined. **Shaanxi** (IZCAS): 2♂, Ningshan, Huoditang, 1538 m, 11–15.VII.2012, leg. Yang Xiushuai, Cheng Rui; 1♂, same locality, 1520 m, 13–17.VIII.2016, leg. Cheng Rui, Jiang Shan; 4♂5♀, same locality, 1497 m, 29–31.VII.2018, leg. Zhang Xinyi; 2♂, Baoji, Jialingjiangyantou, 1620 m, 8–9.VII.2014, leg. Xue Dayong, Ban Xiaoshuang; 2♂1♀, Nanzheng, Liping, 1540 m, 27–30.VII.2017, leg. Li Henan; 1♂, Zhashui, Yingpanzhen, Niubeiliang, Laolin, 1046 m, 11–15.VII.2017, leg. Cui Le; 8♂7♀, Zhashui, Yingpanzhen, Niubeiliang, 1373 m, 24–26.VII.2018, leg. Zhang Xinyi; 1♂, Foping, Longcaoping, 1218 m, 4.VIII.2018, leg. Zhang Xinyi. **Hubei** (IZCAS): 1♂, Yichang, Dengcunxiang, Dalaoling; 1709 m, 15.VII.2013, leg. Cheng Rui. **Sichuan** (IZCAS): 1♀, Emei Shan, Jiulaodong; 1800–1900 m, 28.VII.1957, leg. Huang Keren; 1♂, Baoxing, Dashuigou, 1591 m, 1–5.VIII.2016, leg. Cui Le; 1♂1♀, Jiguan Shan, Shaoyaogou, 1556 m, 11–16.VII.2016, leg. Cui Le. **Yunnan** (IZCAS): 4♂, Pingbian, Dawei Shan, 2090 m, 4–8.VIII.2017, leg. Cui Le.

**Distribution.** China (Shaanxi, Zhejiang, Hubei, Fujian, Sichuan, Yunnan).



**Figures 152–160.** Female genitalia of *Ditrigona* **152** *D. derocina* **153** *D. diana* **154** *D. crystallina*, allotype **155** *D. spilota*, ZFMK **156** *D. furvicosta* **157** *D. sericea* **158** *D. quinaria erminea*, ZFMK **159** *D. obliquilinea thibetaria* **160** *D. uniuncusa*. Scale bars: 1 mm.

**39. *Ditrigona clavata* Li & Wang, 2015**

Figs 44, 80, 114, 147, 176

*Ditrigona clavata* Li & Wang, 2015: 567. Holotype ♂, China: Guangxi, Mao'ershan National Nature Reserve (SCAU).

**Material examined.** CHINA: Shaanxi (IZCAS): 1♂, Liuba, Chengguanzhen, 1007 m, 21–22.VI.2012, leg. Li Jing; 1♂, same locality, 966 m, 23.VI.2012, leg. Liu Shuxian; 1♂, Yang Xian, Huayangzhen, 1099–1108 m, 25–27.VI.2012, leg. Li Jing; 7♂1♀, Ningshan, Yueba, 1052 m, 1–3.VIII.2018, leg. Zhang Xinyi; 2♂, Kang Xian, Qinghelinchang, 1400 m, 8.VII.1999, leg. Zhu Chaodong. Guangxi (IZCAS): 2♂1♀, Huanjiang, Yangmeiao, 1189 m, 18–22.VII.2015, leg. Jiang Nan, Li Xinxin.

**Distribution.** China (Shaanxi, Gansu, Guangdong, Guangxi).

**40. *Ditrigona marmorea* Wilkinson, 1968**

Fig. 45

*Ditrigona marmorea* Wilkinson, 1968: 471. Holotype ♂, Assam: Mishmi Hills (NHMUK).

**Material examined.** CHINA: Yunnan (ZFMK): 1♂, paratype, Li-kiang (China), Provinz Nord-Yuennan, 25.VI. 1935, H. Höne, moth photograph examined.

**Distribution.** China (Yunnan), India.

**41. *Ditrigona quinquelineata* (Leech, 1898)**

*Leucodrepana quinquelineata* Leech, 1898: 364. Holotype ♂, Japan (NHMUK).

*Auzatella quinquelineata*: Inoue, 1962: 13.

*Ditrigona quinquelineata*: Wilkinson, 1968: 480.

**Material examined.** No.

**Distribution.** China (Sichuan), Japan.


**42. *Ditrigona aphyra* Wilkinson, 1968**

Fig. 46

*Ditrigona aphyra* Wilkinson, 1968: 485. Holotype ♂, China: Shaanxi, Tapaishan-im-Tsinling (ZFMK).

**Material examined.** CHINA: Shaanxi (ZFMK): 1♂, holotype, Tapaishan-im-Tsinling Sued-Shensi, ca. 1700 m, 20.VI.1936, H. Höne, moth photograph examined.

**Distribution.** China (Shaanxi).





**Figures 161–172.** Female genitalia of *Ditrigona* **161** *D. tenuiata*, paratype **162** *D. regularis* **163** *D. sinespina* sp. nov., paratype **164** *D. parva* sp. nov., paratype **165** *D. typhodes* **166** *D. lineata lineata* **167** *D. artema* **168** *D. candida*, paratype, ZFMK **169** *D. chionea* **170** *D. fusca* sp. nov., paratype **171** *D. conflexaria micronioides* **172** *D. margarita*. Scale bars: 1 mm.

**43. *Ditrigona cirruncata* Wilkinson, 1968**

Figs 47, 81, 115, 148–151, 177

*Ditrigona cirruncata* Wilkinson, 1968: 497. Holotype ♂, China: Sichuan, Siao-lou (ZFMK).

**Material examined. China: Sichuan:** 1♂ (ZFMK), paratype, Kwanhsien Dist. Suchwan, 16.VIII.1925, leg. G.M. Franck, moth photograph examined; 31♂14♀ (IZCAS), Emei Shan, Qingyinge, 800–1000 m, 14.IV., 16.IV., 17.IV., 18.IV., 19.IV., 24.IV., 25.IV., 26.IV., 27.IV., 29.IV., 30.IV., 1.V., 2.V., 5.V., 6.V., 20.VI., 29.VI., 16.VIII., 18.IX., 20.IX.1957, leg. Wang Zongyuan, Zhu Fuxing, Huang Keren, Lu Youcai; 3♂ (IZCAS), Jiulong Shan, Shizipo, 810 m, 29–31.VII.2016, leg. Cui Le; 2♀, Baoxing, Dashuigou, 1591 m, 1–5.VIII.2016, leg. Cui Le; 1♂2♀ (IZCAS), Hongya, Wawu Shan, Jinhuaqiao, 1147 m, 12–14.VIII.2016, leg. Cui Le; 1♂5♀ (MHBU), Emei Shan, 17–19.IX.2010, leg. Niu Yiping. **Shanxi** (IZCAS): 1♂, Qinshui, Manghe, 557 m, 19–20.VIII.2018, leg. ZhangXinyi. **Henan** (IZCAS): 1♂, Baotianman, 1407 m, 10–11.VIII.2008, leg. Jiang Nan. **Shaanxi:** 11♂17♀ (IZCAS), Ningshan, Huoditang, 1520 m, 13–17.VIII.2016, leg. Cheng Rui, Jiang Shan; 3♂9♀ (IZCAS), same locality, 1497 m, 29–31.VII.2018, leg. Zhang Xinyi; 1♂ (IZCAS), Ningshan, Yueba, 1052 m, 1–3.VIII.2018, leg. Zhang Xinyi; 1♂ (IZCAS), Zhashui, Yingpanzhen, Niubeiliang, 1373 m, 24–26.VII.2018, leg. Zhang Xinyi; 1♀ (IZCAS), Foping, Longcaoping, 1218 m, 4.VIII.2018, leg. Zhang Xinyi; 1♀ (MHBU), Ningshan, Huoditang, 1505 m, 14.VIII.2013, leg. Zhu Xichao, Tian Ying; 1♀ (MHBU), Ningshan, Guanghuojie, 1135 m, 10.VIII.2013, leg. Zhu Xichao, Tian Ying. **Gansu** (IZCAS): 3♀, Wen Xian, VI.–IX.2002, leg. Wang Hongjian; 3♂, Bikou, Bifenggou, 720 m, 8–10.VIII.2016, leg. Cheng Rui, Jiang Shan. **Zhejiang** (IZCAS): 3♂, Lin'an, West Tianmushan, 400 m, 26–27.VII.2003, leg. Xue Dayong, Han Hongxiang; 2♀, West Tianmushan, Xianrending, 1506 m, 27.VII.2011, leg. Yan Keji, Cheng Rui; 1♀, West Tianmushan, Qianmutian, 1330 m, 30.VII.2011, leg. Yan Keji, Cheng Rui; 1♂1♀, Yuyao, Simingshan, 809–853 m, 22–22.VII.2015, leg. Cheng Rui. **Anhui:** 1♀ (MHBU), Shitai, Shanshan, 7.VIII.2010, leg. Ba Yibin, Zhang Zhenxing. **Hubei** (IZCAS): 2♀, Shennongjia, Honghua, 860 m, 21.VIII.1981, leg. Han Yinheng; 1♂, Ying Shan, Taohuachong, 590 m, 23–27.VI.2014, leg. Jiang Nan; 4♂, Xuanen, Changtanhe, Lianghekou, 949 m, 13–14.V.2017, leg. Li Henan; 1♀, Xuanen, Changtanhe, Dawolong, 713 m, 15.V.2017, leg. Li Henan; 2♂1♀, same locality and collector, 794 m, 16.V.2017, leg. Li Henan; 1♀, Lichuan, Xingdou Shan, Sanxianchang, 1144 m, 17–19.V.2017, leg. Li Henan. **Jiangxi** (IZCAS): 1♂, Kuling, 13.VI.1974, leg. Zhang Baolin. **Hunan** (IZCAS): 1♂, Sangzhi, Bamaoxi, Shuitiannan, 370 m, 1.VIII.2009, leg. Wei Zhongmin; 1♂1♀, Sangzhi, Bamaoxi, Shuitianba, 370 m, 5–6.VIII.2009, leg. Wei Zhongmin; 1♂, Sangzhi Badagong Shan, Xiaozhuangping, 1420 m, 18.VI.2015, leg. Yao Jian, Zhao Kaidong; 2♂1♀, Yongshun, Xiaoxixiang, Xiaoxicun, 463–506 m, 21–24.IV.2018, leg. Zhao Kaidong. **Guangxi** (IZCAS): 1♀, Mao'er



178

**Figures 173–178.** Female genitalia of *Ditrigona* **173** *D. berres* **174** *D. chama* **175** *D. platytes* **176** *D. clavata* **177** *D. cirruncata* Scale bars: 1 mm. **178** Neighbour-Joining (NJ) tree of selected *triangularia* species group based on the Kimura two-parameter model.

Shan, Jiuniuchang, 1100 m, 11.VII.1985, leg. Fang Chenglai; 1♂, Mao'er Shan, Jiuniutang, 1146 m, 16.VIII.2012, leg. Cheng Rui; 1♀, Mao'er Shan, Antangping, 1579 m, 17–18.VIII.2012, leg. Cheng Rui.

**Distribution.** China (Shanxi, Henan, Shaanxi, Gansu, Anhui, Zhejiang, Hubei, Jiangxi, Hunan, Guangdong, Guangxi, Sichuan).

**Remarks.** There are some variations in the eighth tergite (Figs 148–151), even in specimens collected from the same locality: for example, both Fig. 149 and Fig. 150 are from Emei Shan, Sichuan Province. Inoue (1962) recorded the distribution of *D. virgo* in central and west China, and was followed by Chu and Wang (1988, 1991). However, when checking the collection of IZCAS, only *D. cirruncata* was found. The record of *D. virgo* in China is doubtful. The situation is similar in *D. komarovi* (Kurentzov), a species recorded from Manchuria, and was combined from *Leucodrepana* by Wilkinson (1968). Chu and Wang (1988, 1991) recorded this species from Northeast China. However, when examining the collection of IZCAS, this species was not found, and its record in China is also doubtful.

## Acknowledgements

We express our sincere thanks to the staff of the Natural History Museum, London, to Dr Dieter Stüning and Dr Marianne Espeland (ZFMK) for allowing examination of material under their curation. We especially thank Sir Anthony Galsworthy, London, for providing the figures of *D. triangularia* and *D. sericea*, correcting the English, and giving valuable suggestions for the latin names of the new species. Thanks also to Dr. Nan Jiang (IZCAS), for taking photographs of type materials of ZFMK. This project was supported by the National Science Foundation of China (31872967, 32170464), the National Science & Technology Fundamental Resources Investigation Program of China (Grant No. 2019FY101800), Biological Resources Programme of Chinese Academy of Sciences (Grant No. KFJ-BRP-017-45), and by a grant (Y229YX5105) from the Key Laboratory of the Zoological Systematics and Evolution of the Chinese Academy of Sciences.

## References

- Ban XS, Jiang N, Cheng R, Xue DY, Han HX (2018) Tribal classification and phylogeny of Geometrinae (Lepidoptera: Geometridae) inferred from seven gene regions. *Zoological Journal of the Linnean Society* 184(3): 653–672. <https://doi.org/10.1093/zoolinnea/zly013>
- Bryk F (1943) Entomological results from the Swedish expedition 1934 to Burma and British India. *Lepidoptera: Drepanidae*. *Arkiv för Zoologi* 34A(13): 1–30.
- Chu HF, Wang LY (1987) A successive Report on the Chinese Drepaninae (Lepidoptera: Drepanidae) Genera: *Albara*, *Auzatella*, *Paralbara*, *Strepsigonia*, *Deroca*, *Cilix* and *Pseudalbara*. *Sinozoologia* 5(5): 105–122.

- Chu HF, Wang LY (1988) On the Chinese species of the genus *Ditrigona* Moore (Lepidoptera: Drepanidae). *Sinozoologia* 6: 199–208.
- Chu HF, Wang LY (1991) *Fauna Sinica, Insecta, Vol. 3, Lepidoptera, Cyclidiidae, Drepanidae*. Beijing, Science Press, 269 pp.
- Dudgeon GC (1899) A catalogue of the Heterocera of Sikkim and Bhutan. Part 6. *Journal of the Bombay Natural History Society* 12: 643–658.
- Gaede M (1931) Family: Drepanidae. In: Strand E (Ed.) *Lepidopterorum Catalogus*, 49, Berlin, 1–60.
- Hampson GF (1893) *The Fauna of British India including Ceylon and Myanmar. Moths 1*. Taylor & Francis, London, 526 pp.
- Hampson GF (1912) The moths of India. Supplementary paper to the Volumes in “The Fauna of British India”. Series IV. Part 6. *Journal of the Bombay Natural History Society* 21: 1222–1272.
- Holloway JD (1998) The moths of Borneo: Families Castniidae, Callidulidae, Drepanidae and Uraniidae. *Malayan Nature Journal* 52: 20–22.
- Inoue H (1962) *Insecta Japonica 2*. Hokuryukan Publishing, Tokyo, 54 pp.
- Jiang S, Han HX (2019) A new species of *Ditrigona* Moore (Lepidoptera: Drepanidae) in China, with a checklist to the *triangularia*-group. *Entomotaxonomia* 41(2): 81–88.
- Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16(2): 111–120. <https://doi.org/10.1007/BF01731581>
- Leech JH (1898) Lepidoptera Heterocera from Northern China, Japan and Korea. *Transactions of the Royal Entomological Society of London* 15: 261–379. <https://doi.org/10.5962/bhl.title.22195>
- Li Y, Xin DY, Wang M (2015) A New Species of the Genus *Ditrigona* Moore, 1888 (Lepidoptera: Drepanidae) in China. *Florida Entomological Society* 98(2): 567–569. <https://doi.org/10.1653/024.098.0227>
- Moore F (1867) On the Lepidopterous Insects of Bengal. *Proceedings of the Zoological Society of London* 1867: 44–98, 612–686.
- Moore F (1888) Heterocera Continued (Pylalidae, Crambidae, Geometridae, Tortricidae, Tineidae). In: Hewitson WC, Moore F (Eds) *Descriptions of New Indian Lepidopterous Insects from the collection of the late Mr. W.S. Atkinson (3)*. Asiatic Society of Bengal, Calcutta, 199–299.
- Oberthür C (1923) Révision iconographique des espèces de Phalénites (*Geometra* Linné) énumérées et décrites par Guenée dans le volume X du *Species général des Lépidoptères*, publié à Paris, chez l’éditeur Roret en 1857. *Études de Lépidoptérologie Comparée* 20: 214–283.
- Poujade GA (1895) Nouvelles espèces de Lépidoptères Hétérocères. *Annales de la Société Entomologique de France* 64: 307–316.
- Saitou N, Nei M (1987) The neighbor-joining method, A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4: 406–425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>
- Strand EH (1917) H. Sauter’s Formosa-ausbeute: Lithosiinae, Nolinae, Noctuidae (p.p.), Ratardidae, Chalcosiidae, sowie nacräge zu den familien Drepanidae, Limacodidae,

- Gelechiidae, Oecophoriidae und Heliodinidae. Archiv für Naturgeschichte 82(A)3: 111–152. <https://doi.org/10.5962/bhl.part.25112>
- Strand E (1910–1911) 12. Family: Drepanidae. In: Seitz A (Ed.) The Macrolepidoptera of the World. Vol. 2: the Palearctic Bombyces & Sphinges. Alfred Kernen, Stuttgart, 195–206.
- Walker F (1861) List of Specimens of Lepidopterous Insects in the Collection of the British Museum, part 23, Edward Newman, London, 757–1020.
- Warren W (1922–1928) Family: Drepanidae. In: Seitz A (Ed.) The Macrolepidoptera of the World. Vol. 10. Bombyces and Sphinges of the Indo-Australian Region. Verlag A. Kernen, Stuttgart, 443–490.
- Watson A (1959) A Revision of the genus *Auzata* Walker (Lepidoptera, Drepanidae). Bonner Zoologische Beiträge 9: 232–257.
- Wilkinson C (1968) A Taxonomic revision of the genus *Ditrigona* (Lepidoptera: Drepanidae: Drepaninae). Transactions of the Zoological Society of London 31(4): 407–517. <https://doi.org/10.1111/j.1096-3642.1968.tb00369.x>