

多曲杯形珊瑚 *Pocillopora meandrina* Dana, 1846 (图版II 图5; 彩图4)

Pocillopora meandrina Dana, Vaughan, 1907, U. S. Nat. Mus. Bull., p. 97, pl. 14, figs. 3, 4; pl. 22, figs. 1, 1a, 2, 2a; pl. 23.

Pocillopora meandrina Dana, Vaughan, 1918, Dept. Mar. Biol. Pub. Carnegie Inst. Wash., p. 78.

Pocillopora meandrina Dana, Yabe, Sugiyama et Eguchi, 1936, Sci. Rep. Tohoku Imp. Univ., 2nd (Geol.), 1, p. 14, pl. 4, fig. 1.

Pocillopora meandrina Dana, Nemenzo, 1964, Nat. Appl. Sci. Bull., p. 215, pl. 10, fig. 1.

Pocillopora meandrina Dana, Scheer & Pillai, 1974, Zoologica, 122, p. 14.

Pocillopora meandrina Dana, Veron & Pichon, 1982, Aust. Inst. Sci. Monogr., Ser. 4, p. 133, figs. 274—276.

Pocillopora meandrina Dana, Dai, 1989, Acta Oceanogr. Taiwanica, p. 93, figs. 13, 17.

Pocillopora meandrina Dana, Veron, 1993, Univ. Hawaii Press, p. 76, figs. 1—3.

Pocillopora meandrina nobilis Verrill, 邹仁林, 1978, 科学出版社, p. 93, pl. 2, figs. 1, 2.

标本采集地: 西沙群岛甘泉岛、晋卿岛、中建岛、盘石屿、永兴岛、南沙群岛美济礁、仁爱礁、信义礁、半月礁、海口礁。

特征: 群体大小不一致的分枝组成, 该种的重要特征是有蔓延的分枝或小枝, 其珊瑚疣较小。隔片和轴柱一般无或发育不全。

生活时为褐色、粉红色或淡黄色。

地理分布: 马尔代夫, 尼科巴群岛, 斑达海, 菲律宾, 塔希堤, 大堡礁, 帛琉群岛, 夏威夷, 琉球群岛, 小笠原群岛; 我国台湾南部、西沙群岛和南沙群岛。

埃氏杯形珊瑚 *Pocillopora eydouxi* (Milne-Edwards & Haime), 1860 (彩图5)

Pocillopora eydouxi (Milne-Edwards & Haime), Vaughan, 1918, Dept. Mar. Biol. Pub. Carnegie Inst. Wash., p. 79, pl. 24, figs. 1—2a.

Pocillopora eydouxi (Milne-Edwards & Haime), Yabe, Sugiyama & Eguchi, 1936, Sci. Rep. Tohoku Imp. Univ., 2nd Ser. (Geol.), Special 1, p. 13, pl. 2, fig. 5; pl. 6, figs. 4, 5; pl. 7, figs. 4—6.

Pocillopora eydouxi (Milne-Edwards & Haime), Umbgrove, 1939, Zool. Med. Mus. Leiden, p. 122; 1940, p. 273, pl. 21, figs. 3, 4, 8.

Pocillopora eydouxi (Milne-Edwards & Haime), Crossland, 1952, Sic. Rep. Great Barrier Reef Exped., 6 (3), p. 112, pl. 1, fig. 2.

Pocillopora eydouxi (Milne-Edwards & Haime), Wells, 1954, U. S. Geol. Surv. Prof., Paper 260-I, p. 414, pl. 98, figs. 3, 4.

Pocillopora eydouxi (Milne-Edwards & Haime), Scheer & Pillai, 1974, Zoologica, 122, p. 15, pl.