



图 102 浦地夜蛾 *Agrotis ripae* (Hubner) ♂ 外生殖器

奇地夜蛾 *Agrotis eugramma* Hampson (图版 IX-1)

Agrotis eugramma Hampson, 1903, Cat. Lep. Phal. Br. Mus. 4: 432, pl. 71, fig. 11. (四川康定)

Feltia miranda Corti, 1928, Mitt. munch. ent. Ges. 18: 13, fig. 3.

Rhyacia eugramma: Warren, 1909, in Seltz, Macrolep. World 3: 47, pl. 10g.

Agrotis eugramma: Corti, 1933, in Seltz, Macrolep. World 3 Suppl. p. 50.

翅展 35 mm。头部与胸部暗褐色，鳞、毛端部灰色；前翅黄褐色带有黑棕色，基线仅在前缘区及中室可见黑色粗点状，内线双线黑色，外一线粗，微波浪形外斜，剑纹小，褐色黑边，环纹圆形，浅褐色，黑边，肾纹黑棕色，黑边，中线模糊，暗褐色，仅中室后方可见，外线双线黑色，细锯齿形，外一线弱，自前缘脉后外弯，至 4 脉后内斜，亚端线黄灰色，稍有间断，外凸于 7 脉，2—5 脉间外弯，内侧有一列锯齿形黑纹，端线由一系列黑点组成；后翅浅赭黄色微带褐色，隐约可见褐色外线，端线褐色，缘毛黄白色，中有一褐线腹部暗赭黄色杂有灰色。

分布 四川。

褐地夜蛾 *Agrotis indiana* Guenée (图版 IX-2)

Agrotis indiana Guenée, 1852, in Boisduval & Guenée, Noct, i: 302, (Indes Orientalis)

Tiracola violacea Butler, 1889, Ill. Het. 7: 54.

Hypernaenia violacea: Hampson, 1894, Fn. Br. Ind. 2: 196.

Mythimna indiana: Hampson, 1903, Cat. Lep. Phal. Br. Mus. 4: 608.