

BMNH: 1♂, lectotype, Khasi Hills, 94-66 (with red type label; original identification label: *Herochroma baba* ♂, Swinhoe, type); 3♂♂, Khasia Hills, Assam, Rothschild Bequest, B. M. 1939-1 (BMNH slide NO. 8495); 1♂, Khasia Nat. Coll., Collection H. J. Elwes, Rothschild Bequest, B. M. 1939-1; 1♂, Sikkim, 1300 ft, Aug. 1897; 1♂, labeled as 'PAHANG, F. M. S. Cameron, Highlands, 4800 ft, 18. 6. 1935, *Herochroma baba latinifrata* Prout type, 2460; with red type label'.

ZFMK: 1♂, Kwangtung: Lianping, 1922. V. 19, H. Höne.

**分布:** 湖北、湖南、福建、广东、广西；印度东北部，尼泊尔，越南，马来半岛。

### (33) 巴始青尺蛾 *Herochroma baibarana* (Matsumura, 1931)

*Dindica baibarana* Matsumura, 1931, 6000 illust. *Insects Japan-Empire*: 893, figured. Holotype ♂, Taiwan: Nantou Hsien, Puli.

*Archaeobalis cristata* Holloway, 1976, *Moths of Borneo with special reference to Mount Kinabalu*: 59, fig. 395, (nec Warren, 1894).

*Archaeobalis orientalis* Holloway, 1982, in Barlow, *An Introduction to the moths of South East Asia*: 242.

*Archaeobalis baibarana*: Inoue, 1986b, *Bull. Fac. domest. Sci.*, *Otsuma Wom. Univ.*, 22: 212, figs 1, 2.

*Herochroma orientalis*: Holloway, 1996, *Malay. Nat. J.*, 49 (3-4): 199, fig. 114, pl. 5: 2.

*Herochroma baibarana*: Inoue, 1999, *Tinea*, 16 (2): 90. figs 23—25.

额具红色带。♂后足胫节有端突。♂第3腹节腹板中部具1对接合的刚毛斑。

翅面暗绿色。前后翅均有黑色小中点。前翅外缘波曲；内线有2个大齿，上方齿较尖，下方齿较圆；外线锯齿形，外伴有浅色阴影及棕褐色杂黑褐色斑块，并在CuA<sub>1</sub>下内弯；亚缘线为脉间一系列浅色点；缘线为脉间黑点。后翅外缘锯齿形，外线、亚缘线、缘线基本同前翅；翅基部有红斑。前后翅反面中点圆点状。

♂外生殖器（据 Inoue, 1999）：钩形突二分裂。背兜侧突基半部略细于端半部。颚形突尖。抱器腹强凸起，抱器瓣外缘在抱器背和抱器腹间深凹陷。横带片具1对三角形骨片，尖端尖。囊形突扁平。味刷发达。阳茎细长，阳茎鞘具1细长刺状突。

♀外生殖器：未知。

**检视标本：** BMNH: 1♂, holotype, Taiwan: Nantou Hsien, Puli. (照片)。

ZSM: 1♂, SABAH (Borneo), Route Keingsu Kimanis PK 26, 1300 m, 7. VIII. 1991, J. Haxaira; 1♂, Thailand, Chiang Mai, Doi Inthanon, 24. VIII. 1998, R. P. Pintarana; 1♂, S. Vietnam, Mt. Ngoc Linh, 15°02'N, 107°59'E, 10/25. VIII. 96, Siniav/Afonin.

**分布:** 中国台湾；印度东北部，越南，泰国，斯里兰卡，马来西亚，印度尼西亚。

(34) 斑始青尺蛾 *Herochroma crassipunctata* (Alphéraky, 1888) (图版IV: 3, 4)

*Gnophos crassipunctata* Alphéraky, 1888, Stettin. ent. Ztg., 49: 68. Lectotype♂, Askai. (ZIS)

*Archaeobalis crassipunctata*: Prout, 1912: in Wytsman, Genera Insectorum, 129: 25.

*Archaeobalis sordida* Wehrli, 1928, Int. ent. Z., 21 (46): 455. Holotype♂, [China]: East Turkestan, Yarkand, Mustagata.

头顶浅橄榄灰色。胸腹部背面颜色同翅面。腹部背面有立毛簇。其他特征未知。

前翅长: ♂23—25 mm。翅浅灰色，散布褐色鳞片。前翅内线黑褐色，细弱波状；外线黑褐色，锯齿形，在脉间常断离，其外侧伴有1灰白色带；亚缘线模糊，为脉间褐色小斑块，缘线清晰，在翅脉端有间断。后翅外线、亚缘线和缘线同前翅。前后翅中点大，灰黑色。翅反面：灰白色散布褐色鳞片；中点大而圆、黑色，较正面清晰。

♂外生殖器（据 Inoue, 1999）：钩形突短小，近三角形。背兜侧突细长，杆状。抱器瓣端部圆，抱器背基部凸出，抱器瓣近端部1/3处向前端延伸成三角形骨化带；抱器腹具小端突。横带片具1对骨化突。囊形突略凸出。阳茎中部具1尖齿。

♀外生殖器（据 Inoue, 1999）：后阴片边缘不清晰，褶皱、骨化；前阴片为1对侧骨化片。囊导管很短，骨化、褶皱。囊体长，头端较细，近中部具1小袋状突。

**检视标本：**ZFMK: 1♂, Tadzhikistan, 1978, L, F1. Vantsh, Gumajak, 2000 m, 5-7. VI, Raitviir, Metsaviir & Viidalepp leg, Museum A. Koenig; 1♂, holotype of *Archaeobalis sordida*, Mustagata, Yarkend.

**分布：**新疆；土耳其斯坦，塔吉克斯坦，阿富汗。

(35) 冠始青尺蛾 *Herochroma cristata* (Warren, 1894)

*Actenochroma cristata* Warren, 1894a, Novit. zool., 1: 381. Lectotype♂, Bhotan [Bhutan]; [India]: Sikkim; Celebes (south) [Sulawesi]. (BMNH)

*Actenochroma subopalina* Warren, 1894a, Novit. zool., 1: 382.

*Pseudoterpnia subtepens* (part) Hampson, 1895b, Fauna Br. India (Moths), 3: 479. (nec Walker, 1860)

*Archaeobalis subtepens* (part) Prout, 1912, in Wytsman, Genera Insectorum, 129: 25. (nec Walker, 1860)

*Archaeobalis cristata*: Prout, 1930a, Bull. Hill Mus. Witley, 4: 127.

*Archaeobalis cristata* ab. *subopalina*: Prout, 1932, in Seitz, Macrolepid. World, 12: 46.

*Herochroma cristata*: Holloway, 1996, Malay. Nat. J., 49 (3—4): 199.

**分布：**台湾、海南、广西、四川；印度，不丹，尼泊尔，越南北部，泰国，印度尼