

SYSU: 1 ♀, Canton, South China, Honam Island, P'an-yu District, June 18, 1936, C. Ng.

BMNH: 1 ♂, holotype, Ceylon, 52.62 (green type label; Specimen photog for CHECKLIST AUST LEP, film 80/17; length of forewing 15 mm); 1 ♂, Ceylan [Ceylon], acq Doncaster, 1892, Ex. Oberthür Coll. Brit. Mus. 1927-3; 1 ♂, syntype of *M. coelataria trychera*, Khasis, Feb. 1894, Nat. Coll., Rothschild Bequest B. M. 1939-1 (*Maxates coelataria trychera* Prout ♂ type; red type label); 1 ♂, same locality and date; 1 ♂, Khasia Hills, Assam, Nissary, Det. by L. B. Prout, Joicey Bequest, Brit. Mus. 1934-120; 1 ♂, Burma, Det. by L. B. Prout, Joicey Bequest Brit. Mus. 1934-120; 1 ♀, Taipinshi, Hainan, June 1906, Rothschild Bequest B. M. 1939-1; 2 ♂♂, Youboi, Hainan, June 1904, Rothschild Bequest B. M. 1939-1; 1 ♂, Sikkim, 1-4000 ft, Möller, Collectio H. J. Elwes, Rothschild Bequest B. M. 1939-1; 1 ♀, Sintang, Borneo, inland of Pontianak, Jan. 1910 (Dr. Martin), Rothschild Bequest B. M. 1939-1 (figured specimen, Moths of Borneo, Part 9); 1 ♂, Pulo Laut, Doherty, 96-34 (BMNH slide NO. 16796) (length of forewing 16—19 mm); 1 ♀, S. India, T. R. Bell, B. M. 1934-394 (BMNH slide NO. 17435); 1 ♀, Trevandrum, Travancore, 93-144, Sept. 1893. 1 ♀, Singapore, H. Roberts Coll. B. M. 1926-395.

ZFMK: 1 ♂, paratype, Khasis, Feb. 1894, Nat. Coll.

分布: 广东、海南; 印度, 缅甸, 越南, 斯里兰卡, 马来西亚, 新加坡, 印度尼西亚 (加里曼丹岛)。

(279) 小青尖尾尺蛾 *Maxates dissimulata* (Walker, 1861) (图 318, 图 638, 图版 XVII: 32)

Thalassodes dissimulata Walker, 1861, *List Specimens lepid. Insects Colln Br. Mus.*, **22**: 551. Holotype ♂, Hindostan [India]. (BMNH)

Thaleura marginata Warren, 1894a, *Novit. zool.*, **1**: 392. Holotype ♂, Bhutan. (BMNH)

Gelasma dissimulata: Prout, 1912, in Wytsman, *Genera Insectorum*, **129**: 147.

Gelasma semiprotrusa Inoue, 1989, *Bull. Fac. domest. Sci., Otsuma Wom. Univ.*, **25**: 266, figs. 31, 37. Holotype ♂, Taiwan: Nantou Hsien, Lushan Spa. **Syn. nov.** (BMNH)

Maxates dissimulata: Holloway, 1996, *Malay. Nat. J.*, **49** (3—4): 274.

♂触角基部 2/3 双栉形, 外侧栉齿长于内侧, 最长栉齿约为触角干直径的 4 倍。额黑褐色, 鳞片粗糙。下唇须背面黑褐色, 腹面白色, 尖端伸出额外。头顶大部分白色, 后缘灰绿色。胸腹部背面灰绿色。♂后足胫节膨大, 有毛束和短端突。

前翅长：15—17 mm。翅型和隐纹尖尾尺蛾 *M. quadripunctata* 相似，但后翅尾突较短。前翅外缘直，臀角方；后翅顶角凸出，近方形。翅正面深灰绿色，带黄绿色调，杂有白色鳞片。前翅前缘淡赭黄色，散布有灰色斑点；白色线纹很弱，伴有深绿色；前翅内线波曲；前后翅外线微弱锯齿形；中点为比翅色略深的短条；缘线颜色略深；缘毛深灰色。翅反面浅绿色，隐约可见翅正面的外线；缘线黑色，在翅脉上有间断。

♂外生殖器：钩形突粗壮，尖端急缩为1点。背兜侧突宽大锤状，短于钩形突。颚形突中突扁宽，端部渐细。抱器瓣宽阔，大部分膜质；抱器背微凸出；抱器瓣中部骨化，在抱器背下方有1列由大小不等的不连续的弱骨化突组成的脊，每个突上均具刚毛；抱器腹中部凸出，抱器腹端突极发达，基部骨化较弱，弯曲，端部强骨化，为伸达抱器瓣端部的大尖齿，尖齿上密布微刺；抱器瓣腹缘近端部处略凸出，但不形成耳状突。横带片膜质，中部凸出。阳端基环狭长，中间近“V”字形凹陷。囊形突狭小。具味刷。阳茎细长针状，弯曲。

♀外生殖器：未知。

检视标本：BMNH: 1♂, holotype, E. India (BMNH slide NO. 8663; with green type label); 1♂, holotype of *Thaleura marginata*, Bhutan, Rothschild Bequest, B. M. 1939-1; 1♂, holotype of *Gelasma semiprotrusa*, Lushan Spa, 1200 m (Nantou Hsien) [Taiwan], Aug. 24—27. 1983, H. Yoshimoto leg, Inoue Coll. B. M. 1992-71 (BMNH slide NO. 16564; Slide No. KY-904, 1987, ♂; slide 13542; Type status verified K. Buckmaster 1993; Type photographed in colour).

分布：上海、台湾；日本，印度，不丹，缅甸，斯里兰卡。

(280) 斜尖尾尺蛾 *Maxates dysgenes* (Prout, 1916) (图 319, 图 639, 图版 XVII: 33)

Gelasma dysgenes Prout, 1916a, *Novit. zool.*, **23**: 13. Holotype ♂, [China]: Tibet, Vriana-tong. (BMNH)

Maxates dysgenes: Holloway, 1996, *Malay. Nat. J.*, **49** (3—4): 274.

♂触角双栉形，外侧栉齿长于内侧，尖端1/3线形；♀线形。额及下唇须均黑褐色，鳞片粗糙，♂♀下唇须尖端伸出额外。头顶前半部白色，后半部灰色。胸腹部背面灰色。♂后足胫节膨大，有毛束，2对距。♂第3腹节腹板具1对刚毛斑。

前翅长：♂♀21—22 mm。前翅顶角尖，外缘近平直；后翅外缘尾突小，折角状。翅灰绿色。前翅前缘黄褐色；内外线白色锯齿形；缘毛灰绿色。后翅外线白色锯齿形，中部外凸；缘毛灰绿色，端半部白色。翅反面灰白，无斑纹。

♂外生殖器：钩形突强骨化，端部较钝。背兜侧突膜质，短于钩形突。颚形突中突扁舌状。抱器瓣狭长，腹缘耳状突发达，其下凹陷较深；抱器腹短，端突钝圆球状，具浓密刚毛。横带片不特化。囊形突狭长，端部平截。无味刷。阳茎极纤细，端部骨化。