

分布 云南；日本，印度，印度尼西亚，萨摩亚群岛，夏威夷，新喀里多尼亚。

暗纹纷夜蛾 *Polydesma scriptilis* Guenée (图版 LIX-13)

Polydesma scriptilis Guenée, 1852, in Boisduval et Guenée, Hist. nat. Ins. Noct. 2: 442, pl. 15, fig. 4. (East Indies: Silhet)

Polydesma otiosa Guenée, 1852, in Boisduval et Guenée, Hist. nat. Ins. Noct. 2: 442.

Polydesma scriptilis: Warren, 1913, in Seitz, Macrolep. World 3: 367.

翅展 40 mm。全体灰褐色；前翅密布细黑点，基线、内线、中线及外线黑色，基线、中线及外线波浪形，内线、亚端线锯齿形，后者双线暗褐色，环纹为小黑环，肾纹有二黑点，亚端线外一线前段似斑条；后翅密布细黑点，各横线黑色波浪形，外线与亚端线均双线，翅外缘一列黑点。

分布 江西、海南、广西、贵州、云南；印度，孟加拉国。

蟠夜蛾属 *Pandesma* Guenée

Pandesma Guenée, 1852, in Boisduval et Guenée, Hist. nat. Ins., Noct. 2: 438.

Type-species: *Pandesma quenavadi* Guenée.

Thria Walker, 1858, List Spec. Lep. Colln Br. Mus. 13: 1079.

Certia Walker, 1858, List Spec. Lep. Colln Br. Mus. 14: 1343.

Michera Walker, 1865, List Spec. Lep. Colln Br. Mus. 33: 810.

Vapara Moore, 1882, Descr. new Indian Lep. Ins. Colln late Atk. p. 163.

Subpandesma Berio, 1966, Boll. Soc. ent. ital. 96: 140.

喙发达，下唇须向上伸，第二节达头顶；胸部被平滑鳞片；腹部基节有毛簇；前翅顶角稍尖；雄蛾后翅反面的基部有平滑鳞片。

蟠夜蛾 *Pandesma quenavadi* Guenée (图版 LIX-14)

Pandesma quenavadi Guenée, 1852, in Boisduval et Guenée, Hist. nat. Ins. Noct. 2: 438, pl. 15, fig. 3. (Bengal)

Pandesma quenavadi: Moore, 1884—1887, Lep. Ceyl. 3: 91. —Warren, 1913, in Seitz, Macrolep. World 3: 367. —Chen, 1982, Icon. Het. Sin. 3: 365—Sugi, 1982, in Inoue et al., Moths Jap. 1: 869.

翅展 49 mm。全体灰色；前翅密布细黑点，基线，内线及外线黑色，内线与外线前段锯齿形，中线黑褐色，前段云状向内扩展，其后波浪形，肾纹小，近长方形，亚端线灰白色锯齿形内斜，内侧一褐带，前宽后窄，线外侧大部带褐色，端线波浪形，黑色间断，内侧一列黑点；后翅白色，翅脉有黑点，后缘区带褐色，亚端区至翅外缘有一褐